

**Feasibility Report  
and  
Final Environmental Impact Statement**

**on**

**Coastal Storm Damage Reduction**

**SURF CITY AND NORTH TOPSAIL BEACH  
NORTH CAROLINA**

**Appendix S -**

**Topsail Beach  
Benthic Community Characterization Survey**




**Appendix S -**

**Topsail Beach**

**Benthic Community Characterization Survey**

The following pages contain the technical memorandum prepared by Dial Cordy and Associates, Inc., Environmental Consultants under contract to the U.S. Army Corps of Engineers, Wilmington District. The technical memorandum describes the benthic study of the Topsail Beach offshore borrow sites. The study also includes 4 additional appendices, which are included in the CDROM version of the Final Feasibility and Final EIS, but are not in the printed version.



# **Technical Memorandum**

## **Topsail Beach Benthic Community Characterization Survey Pender County, NC**

**Final**

**May 2007**

**Prepared for:  
U.S. Army Corps of Engineers  
Wilmington District  
P.O. Box 1890  
Wilmington, NC 28402**

**Prepared by:**



**Topsail Beach Benthic Community  
Characterization Survey  
Pender County, NC**

**Technical Memorandum**

**FINAL**

**May 2007**

**Prepared for:**

**U.S. Army Corps of Engineers  
Wilmington District  
P.O. Box 1890  
Wilmington, NC 28402**

**Prepared by:**

**Dial Cordy and Associates Inc.  
201 North Front Street  
Suite 601  
Wilmington, NC 28401**

## TABLE OF CONTENTS

	Page
LIST OF TABLES .....	iii
LIST OF FIGURES .....	iii
1.0 INTRODUCTION .....	1
2.0 METHODS.....	1
3.0 RESULTS .....	4
3.1    Sediment Grain-size Analysis.....	4
3.2    Faunal Composition.....	6
3.3    Species Richness .....	6
3.4    Diversity Indices.....	6
3.5    Faunal Abundance.....	15
3.6    Biomass .....	15
4.0 DISCUSSION .....	20
5.0 LITERATURE CITED .....	21
Appendix A	Data Quality Control Report
Appendix B	Sediment Analysis Report
Appendix C	Station Data Summary Report
Appendix D	Supporting Data Tables

## LIST OF TABLES

	Page
Table 1. Depths and location coordinates for benthic sampling stations off Topsail Beach, NC.....	4
Table 2. Sediment analysis results for benthic sampling stations off Topsail Beach, NC. ....	5
Table 3. Taxonomic species list for benthic sampling stations off Topsail Beach, NC.....	7
Table 4. Comparison of major taxa occurrences and relative abundances for borrow sites off Topsail Beach, NC.....	10
Table 5. Summary of community parameters for benthic sampling stations off Topsail Beach, NC. ....	13
Table 6. Total abundance and percent composition for taxa collected at borrow sites off Topsail Beach, NC.....	16
Table 7. Mean biomass of major taxa at borrow sites off Topsail Beach, NC. ....	19

## LIST OF FIGURES

	Page
Figure 1. Location of borrow areas off Topsail Beach, NC.....	2
Figure 2. Location of benthic sampling stations off Topsail Beach, NC. ....	3
Figure 3. Relative composition of major taxa for borrow sites off Topsail Beach, NC.....	9
Figure 4. Mean number of species and density within borrow sites off Topsail Beach, NC. .	12
Figure 5. Mean Shannon Wiener index ( $H'$ ) and Pielou Evenness value for borrow sites off Topsail Beach, NC.....	14

## **1.0 INTRODUCTION**


Dial Cordy and Associates Inc. (DC&A) was contracted (Contract DACW W912HN-05-D-0014) by the Wilmington District, Corps of Engineers, to conduct a characterization of the benthic community occurring within six proposed sand borrow sites selected for the proposed Shore Protection Project at Topsail Beach (Figure 1). The initial construction of the berm and dune project, extending along approximately five miles of oceanfront will require 3.2 million cubic yards of beach quality sand. The objectives of the study were to determine through benthic sampling the invertebrate abundance, species composition, and biomass within each of the borrow sites; and to perform a qualitative comparison of survey results to the results of other pertinent benthic studies completed along the coasts of North and South Carolina.

## **2.0 METHODS**

Benthic infaunal samples were collected from three to five random locations within each of the six sand borrow sites shown in Figure 2. Five samples were taken at Borrow Site A, three samples at Borrow Site B, and four samples at each of Borrow Sites C, D, E and F. The borrow areas are located 3.0 to 5.5 miles off the south end of Topsail Island in the Atlantic Ocean. A total of 24 benthic samples were collected, along with one separate sediment sample per station (12 samples) for grain-size analysis. Coordinates for each of the 24 sampling stations are listed in Table 1.

Sampling was first attempted using a standard ponar grab on 13 November 2006; however, adequate bottom penetration was not achieved after 14 attempts within three of the six borrow areas. A Shipek grab ( $0.04\text{ m}^2$ ) was used on November 28<sup>th</sup> and 29th to collect the benthic and sediment samples within each of the six borrow areas. Station positions were recorded using a Trimble GEO XT and real time positions were displayed on-board using a Hypach Navigation System. This allowed the boat operator to select stations throughout each of the borrow areas. To standardize samples, only grabs that collected full samples were saved for analysis. Faunal samples were sieved through a 0.5 mm sieve after collection, preserved in 10% buffered formalin, and subsequently transferred to 50% isoproponal once received by Barry Vittor and Associates Inc., prior to sorting and identification. Samples were sorted and all fauna identified to the lowest practical identification level possible. Wet weight biomass for the major taxa found in each sample was also performed following sample processing.

Grain-size analysis of the 12 sediment samples collected was completed using sieve sizes ranging from 4.75 mm (U.S. Standard Sieve No. 4) to 63 micron (U.S. Standard Sieve 230). A Data Quality Control Report summarizing the field activities for each survey day is provided in Appendix A.



## Legend

  Topsail Beach Borrow Sites



0 2 4 6 8 Miles

### Location of Borrow Sites off Topsail Beach

Topsail Beach Benthic Community Characterization Survey, Pender County, NC

Scale: 1 inch = 2 miles

Drawn By: MR

Date: April 2007


Approved By: SD



DIAL CORDY  
AND ASSOCIATES INC  
Environmental Consultation

J06-0989

Figure 1



### Legend

- ◆ Benthic Sampling Station
- Topsail Beach Borrow Sites

0      6,000      12,000      18,000      24,000      Feet



Location of Benthic Sampling Stations off Topsail Beach

Topsail Beach Benthic Community  
Characterization Survey, Pender County, NC

Scale: 1 inch = 6,000 feet	Drawn By: MR
----------------------------	--------------

Date: April 2007	Approved By: SD
------------------	-----------------

DIAL CORDY AND ASSOCIATES INC	
----------------------------------	--

<i>Environmental Consultation</i>	
-----------------------------------	--

J06-0989

Figure 2

**Table 1. Depths and location coordinates for benthic sampling stations off Topsail Beach, NC.**

Borrow Site Station	Depth (m)	Station Coordinates	
A1	13.7	34°18'16.072"	77°39'47.647"
A2	13.3	34°18'02.486"	77°40'21.868"
A3	13.3	34°18'18.236"	77°39'14.008"
A4	14.2	34°17'54.945"	77°38'53.466"
A5	15.1	34°17'28.093"	77°38'33.163"
B1	12.2	34°18'46.266"	77°38'44.909"
B2	13.5	34°18'41.749"	77°38'24.575"
B3	13.2	34°18'57.558"	77°38'37.370"
C1	14.1	34°16'25.432"	77°36'58.081"
C2	14.9	34°16'06.703"	77°37'23.427"
C3	14.6	34°16'18.018"	77°37'44.459"
C4	15.2	34°15'57.406"	77°38'16.160"
D1	14.2	34°18'25.705"	77°36'07.197"
D2	15.3	34°18'13.213"	77°35'43.565"
D3	14.9	34°18'21.953"	77°35'12.038"
D4	14.3	34°18'36.383"	77°35'29.818"
E1	15.7	34°18'54.333"	77°34'14.820"
E2	15.4	34°18'41.374"	77°34'00.078"
E3	14.7	34°18'43.009"	77°33'37.360"
E4	15.3	34°19'01.135"	77°33'54.375"
F1	15.8	34°19'29.216"	77°32'55.069"
F2	15.4	34°19'37.710"	77°32'59.961"
F3	15.5	34°19'44.872"	77°32'40.239"
F4	14.7	34°19'43.171"	77°32'22.118"

### 3.0 RESULTS

The following results include a review of data pertaining to sediment grain-size analysis, faunal composition, species richness, diversity indices, faunal abundance, and biomass. A sediment analysis report prepared by Dr. Wayne Isfording is provided in Appendix B. Appendix C contains individual data summary reports for each sampling station and Appendix D includes the biomass data for all sampling stations and other supporting tables.

#### 3.1 Sediment Grain-size Analysis

Results of the sediment grain-size analysis are summarized in Table 2 for each sampling station. The mean percent sand for Sites A, B, C, D, E, and F were 99.31%, 98.09%, 99.78%, 99.24%, 99.81%, and 99.84; respectively. The mean percent silt-clay fraction was 0.94%, 1.91%, 0.21%, 0.76%, 0.19%, and 0.16% for Sites A through F; respectively. The highest silt-clay content was recorded for Station B-2, with 5.31%; however, the fine fraction left was considered coarse silt rather than clay. The median particle size and sorting

**Table 2. Sediment analysis results for benthic sampling stations off Topsail Beach, NC.**

Station	% Sand	% Silt + Clay	USACE Description	Median	
	Particle Size (phi)	Sorting Coefficient			
A1	98.73	1.27	Sand	2.515	0.454
A2	99.00	1.00	Sand	2.496	0.470
A3	99.22	0.78	Sand	2.476	0.433
A4	99.72	0.28	Sand	2.470	0.504
A5	99.89	0.11	Sand	2.436	0.497
B1	100.00	0.00	Sand	*	*
B2	94.69	5.31	Sand	2.449	0.609
B3	99.59	0.41	Sand	*	*
C1	99.92	0.08	Sand	2.413	0.488
C2	99.82	0.18	Sand	2.445	0.436
C3	99.78	0.22	Sand	2.465	0.429
C4	99.57	0.43	Sand	2.469	0.415
D1	99.86	0.14	Sand	2.268	0.708
D2	99.84	0.16	Sand	2.412	0.505
D3	99.94	0.06	Sand	2.358	0.643
D4	99.61	0.39	Sand	1.885	0.726
E1	97.57	2.43	Sand	2.344	0.627
E2	99.82	0.18	Sand	2.271	0.693
E3	99.95	0.05	Sand	1.679	0.663
E4	99.82	0.18	Sand	1.798	0.711
F1	99.67	0.33	Sand	2.217	0.714
F2	99.78	0.22	Sand	*	0.934
F3	99.85	0.15	Sand	2.375	0.571
F4	99.91	0.09	Sand	2.339	0.618

\* Insufficient data for any statistical calculations (See Appendix B).

coefficient for Sites A and E varied little between sampling stations (Table 2), while these parameters varied more between sampling stations at Sites C and D. Median particle size (phi) and sorting coefficients for Stations B-1, B-3, and F-2 could not be statistically calculated due to insufficient data (See Appendix B).

### **3.2 Faunal Composition**

A total of 104 taxa were collected from the 24 samples collected within the six proposed borrow sites (Table 3). Of the taxa collected, the dominant faunal groups included polychaetes (43), crustaceans (26), mollusks (24), and echinoderms (6). Major taxa composition within each of the six borrow sites are illustrated in Figure 3 and Table 4. As shown, polychaetes were the major component of the benthos within all borrow sites; except for Site C, where crustaceans were more prevalent. The taxa within Site D were generally distributed evenly between polychaetes, mollusks, and crustaceans. Other, less frequently occurring faunal groups, included rhynchocoela (ribbon worms) that occurred within all five borrow sites, ranging from 2.6% to 4.8% of the total taxa collected; and echinoderms, which occurred within four of the five borrow sites, ranging from 3.8% to 14.3% of the total taxa collected. The total number of taxa collected within all of the samples taken at each borrow site ranged from a low of 21 taxa at Site C (n=4) to a high of 39 taxa at Sites B (n=3), E (n=4), and F (n=4) (Table 4).

Based on the total count of individuals collected per taxa within each borrow site, polychaetes were the dominant taxa at Sites A, B, D, and F. Polychaetes comprised greater than 30% of the total count (Table 4, Figure 3). Mollusks were the dominant taxa at Site E (44.6%) and secondary dominant taxa at Sites A (37.5%), B (15.4%), D (31.7%), and E (44.6%). The distribution of the total number of individuals within all Site F samples were fairly evenly distributed between the three most abundant taxa groups (polychaetes, mollusks, and crustaceans).

### **3.3 Species Richness**

The mean number of species present within the six borrow sites ranged from a low of 7.8 within Site C to a high of 16.7 within Site B. The mean number of species present within Sites A (8.4) and D (8) were similar to Site C, while the mean number of species within Site E (15.8) was more similar to the mean number of species present within Site B. The mean number of species within Site F was 12.3 (Figure 4, Table 5). The total number of taxa collected within Sites C and D was also lower than observed within the other four sites (Table 4).

### **3.4 Diversity Indices**


Diversity indices were calculated for each sample collected within the borrow sites (Table 5). Mean Shannon Wiener indices and Pielou Evenness values for the six borrow sites are illustrated on Figure 5. Diversity values expressed as one of the three diversity indices were consistently higher at Sites B, E, and F. High Evenness ( $J'$ ) and Equitability ( $e$ ) values for samples collected within the six borrow sites indicate that within each station, the abundance was evenly distributed among the taxa collected. Evenness values were greater than 0.90 at all but two of the 24 stations. Mean evenness values for the six sites ranged from 0.91 at Site E to 0.96 at Sites A and D. Variability in the evenness values within each site was highest at Site F (Figure 5).

**Table 3. Taxonomic species list for benthic sampling stations off Topsail Beach, NC.**

<b>ANNELIDA</b>	
CLASS OLIGOCHAETA	
Order TUBIFICIDA	
FAMILY ENCHYTRAEIDAE	
<i>Enchytraeidae (LPIL)</i>	
FAMILY TUBIFICIDAE	
<i>Tubificidae (LPIL)</i>	
CLASS POLYCHAETA	
Order CAPITELLIDA	
FAMILY CAPITELLIDAE	
<i>Capitellidae (LPIL)</i>	
<i>Mediomastus (LPIL)</i>	
<i>Mediomastus californiensis</i>	
FAMILY MALDANIDAE	
<i>Maldanidae (LPIL)</i>	
Order EUNICIDA	
FAMILY DORVILLEIDAE	
<i>Schistomerings pectinata</i>	
FAMILY LUMBRINERIDAE	
<i>Lumbrineris latreilli</i>	
FAMILY OENONIDAE	
<i>Arabella multidentata</i>	
FAMILY ONUPHIDAE	
<i>Onuphidae (LPIL)</i>	
Order OPHELIIDA	
FAMILY OPHELIIDAE	
<i>Opheliidae (LPIL)</i>	
<i>Armandia (LPIL)</i>	
<i>Armandia maculata</i>	
<i>Ophelia denticulata</i>	
Order ORBINIIDA	
FAMILY PARAONIDAE	
<i>Aricidea (LPIL)</i>	
<i>Aricidea catherinae</i>	
<i>Aricidea suecica</i>	
<i>Aricidea wassi</i>	
<i>Cirrophorus (LPIL)</i>	
Order OWENIIDA	
FAMILY OWENIIDAE	
<i>Owenia fusiformis</i>	
Order PHYLLODOCIDA	
FAMILY CHRYSOPETALIDAE	
<i>Bhawania heteroseta</i>	
FAMILY GLYCERIDAE	
<i>Glyceridae (LPIL)</i>	
FAMILY GONIADIDAE	
<i>Goniada littorea</i>	
<i>Goniadides caroliniae</i>	
FAMILY HESIONIDAE	
<i>Microphthalmus (LPIL)</i>	
FAMILY NEPHTYIDAE	
<i>Aglaophamus verrilli</i>	
<i>Nephtys picta</i>	
<i>Nephtys simoni</i>	
FAMILY NEREIDAE	
<i>Nereididae (LPIL)</i>	
<i>Ceratocephale oculata</i>	
FAMILY PHYLLODOCIDAE	
<i>Eumida sanguinea</i>	
FAMILY PILARGIIDAE	
<i>Ancistrosyllis (LPIL)</i>	
<i>Ancistrosyllis hartmanae</i>	
<i>Sigambra tentaculata</i>	
Order SPIONIDA	
FAMILY CHAOPTERIDAE	
<i>Spiochaopterus oculatus</i>	
FAMILY CIRRATULIDAE	
<i>Cirratulidae (LPIL)</i>	
<i>Caulieriella sp. J</i>	
FAMILY MAGELONIDAE	
<i>Magelona (LPIL)</i>	
<i>Magelona papillicornis</i>	
<i>Magelona pettiboneae</i>	
FAMILY SPIONIDAE	
<i>Spionidae (LPIL)</i>	
<i>Parapriopospio pinnata</i>	
<i>Spiophanes bombyx</i>	
Order TEREBELLIDA	
FAMILY PECTINARIIDAE	
<i>Pectinaria gouldii</i>	
FAMILY TEREBELLIDAE	
<i>Terebellidae (LPIL)</i>	
<b>ARTHROPODA</b>	
CLASS MALACOSTRACA	
Order AMPHIPODA	
<i>Amphipoda (LPIL)</i>	
FAMILY AORIDAE	
<i>Rildardanus laminosa</i>	
<i>Unciola serrata</i>	
FAMILY HAUSTORIIDAE	
<i>Acanthohaustorius intermedius</i>	
FAMILY ISAEIDAE	
<i>Microprotopus raneyi</i>	
FAMILY MELITIDAE	
<i>Maera caroliniana</i>	
FAMILY OEDICEROTIDAE	
<i>Oedicerotidae (LPIL)</i>	
<i>Americhelidium americanum</i>	
FAMILY PHOXOCEPHALIDAE	
<i>Metharpinia floridana</i>	
<i>Rhepoxyinius hudsoni</i>	
FAMILY PLATYISCHNOPIDAE	
<i>Eudevenopus honduranus</i>	
FAMILY SYNOPIIDAE	
<i>Metatiron (LPIL)</i>	
<i>Metatiron triocellatus</i>	
Order CUMACEA	
FAMILY DIASTYLIDAE	
<i>Oxyurostylis (LPIL)</i>	
Order DECAPODA	
FAMILY PAGURIDAE	
<i>Pagurus (LPIL)</i>	
FAMILY PASIPHAEIDAE	
<i>Leptochela serratorbita</i>	
FAMILY PINNOTHERIDAE	
<i>Dissodactylus mellitae</i>	
CLASS OSTRACODA	
Order MYODOCOPINA	
FAMILY PHILOMEDIDAE	
<i>Philomedidae (LPIL)</i>	
<i>Harbansus paucichelatus</i>	
FAMILY SARSIELLIIDAE	
<i>Eusarsiella (LPIL)</i>	
<i>Eusarsiella radiicosta</i>	
<i>Eusarsiella texana</i>	

**Table 3. (concluded)**

Order PODOCOPIDA <i>Podocopida (LPIL)</i>	Order NEOGASTROPODA FAMILY OLIVIDAE
<b>BRYOZOA</b> <i>Bryozoa (LPIL)</i>	<i>Olivella dealbata</i>
<b>CHORDATA</b>	<i>Oliva sayana</i>
CLASS LEPTOCARDIA	FAMILY TURRIDAE
Order AMPHOXI	<i>Kurtziella atrostyla</i>
FAMILY BRANCHIOSTOMIDAE	Order PYRAMIDELLOIDA
<i>Branchiostoma (LPIL)</i>	FAMILY PYRAMIDEILLIDAE
<b>CNIDARIA</b>	<i>Odostomia weberi</i>
CLASS HYDROZOA	<i>Turbanilla (LPIL)</i>
<i>Hydrozoa (LPIL)</i>	<b>PHORONIDA</b>
<b>ECHINODERMATA</b>	FAMILY PHORONIDAE
<i>Echinodermata (LPIL)</i>	<i>Phoronis (LPIL)</i>
CLASS ECHINOIDEA	<b>PLATYHELMINTHES</b>
<i>Echinoidea (LPIL)</i>	CLASS TURBELLARIA
Order CLYPEASTEROIDA	<i>Turbellaria (LPIL)</i>
FAMILY MELLITIDAE	<b>RHYNCHOCOELA</b>
<i>Mellitidae (LPIL)</i>	<i>Rhynchocoela (LPIL)</i>
<i>Mellita isometra</i>	<b>SIPUNCULA</b>
CLASS OPHIUROIDEA	FAMILY ASPIDOSIPHONIDAE
<i>Ophiuroidea (LPIL)</i>	<i>Aspidosiphon albus</i>
Order OPHIURIDA	
FAMILY AMPHIURIDAE	
<i>Amphiuridae (LPIL)</i>	
<b>MOLLUSCA</b>	
CLASS BIVALVIA	
<i>Bivalvia (LPIL)</i>	
Order OSTREOIDA	
FAMILY ANOMIIDAE	
<i>Anomia simplex</i>	
Order VENEROIDA	
FAMILY CRASSATELLIDAE	
<i>Crassinella dupliniana</i>	
<i>Crassinella lunulata</i>	
FAMILY LUCINIDAE	
<i>Lucinidae (LPIL)</i>	
<i>Lucina (LPIL)</i>	
<i>Lucina multilineata</i>	
FAMILY MESODESMATIDAE	
<i>Ervilia concentrica</i>	
FAMILY MONTACUTIDAE	
<i>Montacutidae (LPIL)</i>	
FAMILY SEMELIDAE	
<i>Semelidae (LPIL)</i>	
FAMILY TELLINIDAE	
<i>Strigilla mirabilis</i>	
<i>Tellina (LPIL)</i>	
<i>Tellina iris</i>	
CLASS GASTROPODA	
Order CEPHALASPIDEA	
FAMILY SCAPHANDRIDAE	
<i>Akteocina canaliculata</i>	
<i>Cylichna alba</i>	
Order MESOGASTROPODA	
FAMILY CAECIDAE	
<i>Caecum pulchellum</i>	
FAMILY EULIMIDAE	
<i>Strombiformis bilineatus</i>	
FAMILY NATICIDAE	
<i>Tectonatica pusilla</i>	



**Figure 3. Relative composition of major taxa for borrow sites off Topsail Beach, NC.**


**Table 4. Comparison of major taxa occurrences and relative abundances for borrow sites off Topsail Beach, NC.**

Site	Cnidaria	Rhynchoela	Annelida	Mollusca	Platyhelminthes	Arthropoda	Bryozoa	Sipuncula	Phoronida	Echinodermata	Chordata	Total Per Site
<b>Borrow Site A (n=5)</b>												
Total # Taxa	1	1	18	9		4			1			34
Taxa % Total	2.9%	2.9%	52.9%	26.5%		11.8%			2.9%			
Total # Ind.	1	1	27	21		5			1			56
Ind. % Total	1.8%	1.8%	48.2%	37.5%		8.9%			1.8%			
<b>Borrow Site B (n=3)</b>												
Total # Taxa		1	19	6	1	7	1	1	1	2		39
Taxa % Total		2.6%	48.7%	15.4%	2.6%	17.9%	2.6%	2.6%	2.6%	5.1%		
Total # Ind.		3	44	16	4	9	1	1	1	9		88
Ind. % Total		3.4%	50.0%	18.2%	4.5%	10.2%	1.1%	1.1%	1.1%	10.2%		
<b>Borrow Site C (n=4)</b>												
Total # Taxa		1	5	4		8			3			21
Taxa % Total		4.8%	23.8%	19.0%		38.1%			14.3%			
Total # Ind.		1	12	16		20			6			55
Ind. % Total		1.8%	21.8%	29.1%		36.4%			10.9%			


**Table 4. (concluded)**

<b>Site</b>	<b>Cnidaria</b>	<b>Rhynchocoela</b>	<b>Annelida</b>	<b>Mollusca</b>	<b>Platyhelminthes</b>	<b>Arthropoda</b>	<b>Bryozoa</b>	<b>Sipuncula</b>	<b>Phoronida</b>	<b>Echinodermata</b>	<b>Chordata</b>	<b>Total Per Site</b>
<b>Borrow Site D (n=4)</b>												
<b>Total # Taxa</b>		1	9	7		8				1		26
<b>Taxa % Total</b>		3.8%	34.6%	26.9%		30.8%				3.8%		
<b>Total # Ind.</b>		1	15	13		11				1		41
<b>Ind. % Total</b>		2.4%	36.6%	31.7%		26.8%				2.4%		
<b>Borrow Site E (n=4)</b>												
<b>Total # Taxa</b>		1	18	10		7				2	1	39
<b>Taxa % Total</b>		2.6%	46.2%	25.6%		17.9%				5.1%	2.6%	
<b>Total # Ind.</b>		4	31	50		23				2	2	112
<b>Ind. % Total</b>		3.6%	27.7%	44.6%		20.5%				1.8%	1.8%	
<b>Borrow Site F (n=4)</b>												
<b>Total # Taxa</b>		1	18	9	1	7				2	1	39
<b>Taxa % Total</b>		2.6%	46.2%	23.1%	2.6%	17.9%				5.1%	2.6%	
<b>Total # Ind.</b>		3	23	21	1	21				4	3	76
<b>Ind. % Total</b>		3.9%	30.3%	27.6%	1.3%	27.6%				5.3%	3.9%	

**Mean Number of Species ( $\pm 1$  SD)**



**Mean Density ( $\pm 1$  SD)**




**Figure 4. Mean number of species and density within borrow sites off Topsail Beach, NC.**


**Table 5. Summary of community parameters for benthic sampling stations off Topsail Beach, NC.**

Station	Total No. Taxa	Total No. Individuals	Density (nos/m <sup>2</sup> )	H' Shannon (log e)	d Diversity (log 2)	1/S Simpson Diversity	J' Pielou Evenness	D Margalef Richness	e Equitability
A1	6	7	175.0	1.75	2.52	21.00	0.98	2.57	1.33
A2	10	17	425.0	2.18	3.15	13.60	0.95	3.18	1.25
A3	9	10	250.0	2.16	3.12	45.00	0.98	3.47	1.37
A4	9	11	275.0	2.15	3.10	27.50	0.98	3.34	1.34
A5	7	11	275.0	1.80	2.59	9.17	0.92	2.50	1.20
B1	14	25	625.0	2.53	3.65	20.00	0.96	4.04	1.29
B2	9	9	225.0	2.20	3.17		1.00	3.64	1.41
B3	27	51	1275.0	2.98	4.29	19.92	0.90	6.61	1.06
C1	3	5	125.0	1.05	1.52	5.00	0.96	1.24	1.29
C2	9	17	425.0	2.04	2.94	10.46	0.93	2.82	1.20
C3	9	16	400.0	1.98	2.85	8.57	0.90	2.89	1.12
C4	10	17	425.0	2.15	3.10	12.36	0.93	3.18	1.21
D1	4	4	100.0	1.39	2.00		1.00	2.16	1.37
D2	8	10	250.0	1.97	2.85	15.00	0.95	3.04	1.26
D3	4	7	175.0	1.28	1.84	5.25	0.92	1.54	1.22
D4	16	20	500.0	2.67	3.85	31.67	0.96	5.01	1.30
E1	11	15	375.0	2.27	3.27	17.50	0.95	3.69	1.25
E2	13	26	650.0	2.34	3.38	11.21	0.91	3.68	1.14
E3	22	43	1075.0	2.64	3.82	10.38	0.86	5.58	0.92
E4	17	28	700.0	2.58	3.72	14.00	0.91	4.80	1.12
F1	9	9	225.0	2.20	3.17		1.00	3.64	1.41
F2	16	24	600.0	2.64	3.80	23.00	0.95	4.72	1.26
F3	9	14	350.0	2.14	3.09	18.20	0.98	3.03	1.34
F4	15	29	725.0	2.20	3.17	5.88	0.81	4.16	0.85

**Mean Shannon Wiener Index ( $H'$ )**



**Mean Pielou Evenness ( $J$ )**



**Figure 5. Mean Shannon Wiener index ( $H'$ ) and Pielou Evenness value for borrow sites off Topsail Beach, NC.**

### **3.5 Faunal Abundance**

The total count and density of the benthic community collected at each station within the six borrow sites is provided in Table 5. The abundance and density for each species occurrence is provided in Tables D-1 and D-2 (Appendix D). A total of 425 individuals were collected during the study, ranging from a mean density of 256 individuals/m<sup>2</sup> within Site D to 708 individuals/m<sup>2</sup> at Site B (Figure 4). Variability in the density within each borrow site was highest at Site B, which ranged from 225 individuals/m<sup>2</sup> to 1,275 individuals/m<sup>2</sup> (n=3); followed by Site E, where density ranged from 375 individuals/m<sup>2</sup> to 1,075 individuals/m<sup>2</sup> (n=4). Mean density at Sites A, C, and D were all low compared to the mean density at Sites B, E, and F.

The ten most abundant species were: the polychaetes *Magelona papillicornis*, *Goniada littorea*, and *Goniadides carolinae*; the amphipod *Eudovenopus honduranus*; the bivalves *Crassinella lunicata*, *Lucinidae*(LPIL), and *Tellina* (LPIL); the gastropod *Caecum pulchellum*; the echinoderm *Ophiuroidea* (LPIL); and the ribbon worm *Rhynchocoela* (LPIL). These ten species together comprised 47.4% of the benthic infauna collected (Table 6). Other species found within at least two of the sites, and comprising about 2% of the total abundance included: the polychaetes *Onuphidae* (LPIL), *Armandia maculata*, *Bhawania heteroseta*, and *Spionidae* (LPIL); the bivalve *Crassinella dupliniana*; and the gastropods *Acteocina canaliculata* and *Cylichna alba*. Of the balance of species remaining, 79 species each comprised less than 1% of the total abundance.

### **3.6 Biomass**

The biomass for major faunal groups collected at all sampling stations is summarized in Table D-3 of Appendix D, while the mean biomass (g/m<sup>2</sup>) for major faunal groups within each site is provided in Table 7. Mean biomass was lowest at Site A (36.69 g/m<sup>2</sup>) and Site D (41.59 g/m<sup>2</sup>); similar between Sites B (69.53 g/m<sup>2</sup>), Site E (59.08 g/m<sup>2</sup>), and Site F (57.85 g/m<sup>2</sup>); and highest at Site C (182.61 g/m<sup>2</sup>). Site C is skewed higher due to the occurrence of brittle stars (*Ophiuroidea*), which weigh more than other taxa present. Excluding this epibenthic taxa, the mean biomass for Site C would be 44.88 g/m<sup>2</sup>.

**Table 6. Total abundance and percent composition for taxa collected at borrow sites off Topsail Beach, NC.**

Species	Total Count Per Site						% of Total
	A	B	C	D	E	F	
<i>Enchytraeidae (LPIL)</i>		2					0.48
<i>Tubificidae (LPIL)</i>		1					0.24
<i>Capitellidae (LPIL)</i>	1						0.24
<i>Mediomastus (LPIL)</i>		4			1		1.20
<i>Mediomastus californiensis</i>	1						0.24
<i>Maldanidae (LPIL)</i>					1		0.24
<i>Schistomerings pectinata</i>		1					0.24
<i>Lumbrineris latreilli</i>				1		1	0.48
<i>Arabella multidentata</i>					1		0.24
<i>Onuphidae (LPIL)</i>	1	1	2	1		2	1.67
<i>Opheliidae (LPIL)</i>					2	1	0.72
<i>Armandia (LPIL)</i>					1		0.24
<i>Armandia maculata</i>	1	1	3		2		1.67
<i>Ophelia denticulata</i>						1	0.24
<i>Aricidea (LPIL)</i>				1	1		0.48
<i>Aricidea catherinae</i>	1	1					0.48
<i>Aricidea suecica</i>					1		0.24
<i>Aricidea wassi</i>				1	3		0.96
<i>Cirrophorus (LPIL)</i>		1			1		0.48
<i>Owenia fusiformis</i>	1			1			0.48
<i>Bhawania heteroseta</i>		7				1	1.91
<i>Glyceridae (LPIL)</i>	2	1	1		1		1.20
<i>Goniada littorea</i>	1		5	4	4	2	3.83
<i>Goniadides carolinae</i>		9				2	2.63
<i>Microphtalmus (LPIL)</i>						1	0.24
<i>Aglaophamus verrilli</i>	1						0.24
<i>Nephtys picta</i>						2	0.48
<i>Nephtys simoni</i>						1	0.24
<i>Nereididae (LPIL)</i>		1					0.24
<i>Ceratocephale oculata</i>					1	2	0.72
<i>Eumida sanguinea</i>		1					0.24
<i>Ancistrosyllis (LPIL)</i>		1			1		0.48
<i>Ancistrosyllis hartmanae</i>		1			1		0.48
<i>Sigambra tentaculata</i>	1						0.24
<i>Spiochaetopterus oculatus</i>	1		1				0.48
<i>Cirratulidae (LPIL)</i>		2				1	0.72
<i>Caulieriella sp. J</i>	4	1				1	1.44
<i>Magelona (LPIL)</i>	3						0.72

**Table 6. (continued)**

Species	Total Count Per Site						% of Total
	A	B	C	D	E	F	
<i>Magelona papillicornis</i>	3			4	7	2	3.83
<i>Magelona pettiboneae</i>						1	0.24
<i>Spionidae (LPIL)</i>	1	4		1	1	1	1.91
<i>Parapriionospio pinnata</i>	1				1	1	0.72
<i>Spiophanes bombyx</i>				1			0.24
<i>Pectinaria gouldii</i>	1						0.24
<i>Terebellidae (LPIL)</i>		1					0.24
<i>Amphipoda (LPIL)</i>			1	1			0.48
<i>Rildardanus laminosa</i>		1					0.24
<i>Unciola serrata</i>		1					0.24
<i>Acanthohaustorius intermedius</i>						2	0.48
<i>Micropotopus raneyi</i>		2					0.48
<i>Maera caroliniana</i>		2				1	0.72
<i>Oedicerotidae (LPIL)</i>		1					0.24
<i>Americhelidium americanum</i>				1	1		0.48
<i>Metharpinia floridana</i>						1	0.24
<i>Rhepoxyinius hudsoni</i>			2	1	1	1	1.20
<i>Eudevenopus honduranus</i>			10	4	17	14	10.77
<i>Metatiron (LPIL)</i>		1					0.24
<i>Metatiron triocellatus</i>		1					0.24
<i>Oxyurostylis (LPIL)</i>				1	1	1	0.72
<i>Pagurus (LPIL)</i>		1				1	0.48
<i>Leptochela serratorbita</i>							
<i>Dissodactylus mellitae</i>			2				0.48
<i>Philomedidae (LPIL)</i>						1	0.24
<i>Harbansus paucichelatus</i>						1	0.24
<i>Eusarsiella (LPIL)</i>	1						0.24
<i>Eusarsiella radiocosta</i>						1	0.24
<i>Eusarsiella texana</i>	1		2	1	1	1	1.44
<i>Podocopida (LPIL)</i>			1				0.24
<i>Bryozoa (LPIL)</i>		1					0.24
<i>Branchiostoma (LPIL)</i>					2	3	1.20
<i>Hydrozoa (LPIL)</i>	1						0.24
<i>Echinodermata (LPIL)</i>						1	0.24
<i>Echinoidea (LPIL)</i>			2				0.48
<i>Mellitidae (LPIL)</i>						2	0.48
<i>Mellita isometra</i>			1				0.24
<i>Ophiuroidea (LPIL)</i>		8	3	1	1	2	3.59
<i>Amphiuridae (LPIL)</i>		1					0.24
<i>Bivalvia (LPIL)</i>	2	1					0.72

**Table 6. (concluded)**

Species	Total Count Per Site						% of Total
	A	B	C	D	E	F	
<i>Anomia simplex</i>						1	0.24
<i>Crassinella dupliniana</i>					4	4	1.91
<i>Crassinella lunulata</i>		10	2	3			3.59
<i>Lucinidae (LPIL)</i>	2			1	3	3	2.15
<i>Lucina (LPIL)</i>				1			0.24
<i>Lucina multilineata</i>		1			3		0.96
<i>Ervilia concentrica</i>	1			1		1	0.72
<i>Montacutidae (LPIL)</i>					1		0.24
<i>Semelidae (LPIL)</i>		1					0.24
<i>Strigilla mirabilis</i>					2		0.48
<i>Tellina (LPIL)</i>	5	1	9	1	3		4.54
<i>Tellina iris</i>	1						0.24
<i>Acteocina canaliculata</i>	1		1	1	4	1	1.91
<i>Cylichna alba</i>	5		2	1			1.91
<i>Caecum pulchellum</i>		1	4	7	24	3	9.33
<i>Strombiformis bilineatus</i>					2		0.48
<i>Tectonatica pusilla</i>			1		2		0.72
<i>Oliva sayana</i>	3						0.72
<i>Olivella dealbata</i>						2	0.48
<i>Kurtziella atrostyla</i>						1	0.24
<i>Odostomia weberi</i>	1						0.24
<i>Turbanilla (LPIL)</i>						1	0.24
<i>Phoronis (LPIL)</i>	1	1					0.48
<i>Turbellaria (LPIL)</i>		4				1	1.20
<i>Rhynchocoela (LPIL)</i>	1	3	1	1	4	3	3.11
<i>Aspidosiphon albus</i>		1					0.24

**Table 7. Mean biomass of major taxa at borrow sites off Topsail Beach, NC.**

<b>Borrow Site A</b>		<b>Borrow Site D</b>	
<b>Faunal Group</b>	<b>gm/m<sup>2</sup></b>	<b>Faunal Group</b>	<b>gm/m<sup>2</sup></b>
<b>Annelida</b>	14.3835	<b>Annelida</b>	14.1350
<b>Mollusca</b>	14.0410	<b>Mollusca</b>	13.7320
<b>Arthropoda</b>	5.5235	<b>Arthropoda</b>	6.8550
<b>Echinodermata</b>	0.0000	<b>Echinodermat a</b>	3.4325
<b>Other Taxa</b>	2.7450	<b>Other Taxa</b>	3.4325
<b>Total</b>	36.6930	<b>Total</b>	41.5881
<b>Borrow Site B</b>		<b>Borrow Site E</b>	
<b>Faunal Group</b>	<b>gm/m<sup>2</sup></b>	<b>Faunal Group</b>	<b>gm/m<sup>2</sup></b>
<b>Annelida</b>	13.9200	<b>Annelida</b>	14.0706
<b>Mollusca</b>	13.7866	<b>Mollusca</b>	13.8350
<b>Arthropoda</b>	13.9208	<b>Arthropoda</b>	14.0100
<b>Echinodermata</b>	13.9516	<b>Echinodermat a</b>	6.8643
<b>Other Taxa</b>	13.9500	<b>Other Taxa</b>	10.2950
<b>Total</b>	69.5291	<b>Total</b>	59.0750
<b>Borrow Site C</b>		<b>Borrow Site F</b>	
<b>Faunal Group</b>	<b>gm/m<sup>2</sup></b>	<b>Faunal Group</b>	<b>gm/m<sup>2</sup></b>
<b>Annelida</b>	13.8419	<b>Annelida</b>	15.0306
<b>Mollusca</b>	13.7693	<b>Mollusca</b>	13.9231
<b>Arthropoda</b>	13.8356	<b>Arthropoda</b>	13.8325
<b>Echinodermata</b>	137.7275	<b>Echinodermata</b>	3.4350
<b>Other Taxa</b>	3.4350	<b>Other Taxa</b>	11.6306
<b>Total</b>	182.6093	<b>Total</b>	57.8518

## **4.0 DISCUSSION**

In comparison to the results of baseline sampling performed for the Kure Beach restoration project (Posey and Alphin 2000, 2002) and for the Dare County beach shoreline protection project (Versar 2006), the benthic community found offshore of Topsail Beach was less diverse and abundant. Since the Topsail Beach benthic sampling in this survey was only performed in late fall/early winter, comparisons to other surveys should focus on the same seasonal time frame.

Baseline sampling performed off Kure Beach in October 1995 found 44 and 76 taxa, and faunal densities of 1,007 individuals/m<sup>2</sup> and 923 individuals/m<sup>2</sup>, respectively, for the borrow and control sites (Posey and Alphin 2000). The number of total taxa occurring within each proposed borrow site off Topsail Beach ranged from 21 to 39 taxa, while mean density by site ranged from 280 individuals/m<sup>2</sup> to 708 individuals/m<sup>2</sup>. Posey and Alphin (2000) reported 600 total taxa collected from the year long study, as compared to the 104 total taxa collected for the one-time sampling performed for the Topsail Beach study. However, the Kure Beach study was more extensive in that it involved five sampling periods, replicate samples at each location, and covered a variety of benthic substrates (some adjacent to hard bottom communities). All of these factors would be expected to yield a more diverse benthic community. The community studied off Kure Beach was dominated by polychaetes, with 29 taxa comprising at least 5% of the individuals sampled from a single sampling event. Off Topsail Beach, polychaetes dominated the community, comprising over 30% of the relative abundance at four of the six borrow sites.

Seasonal pre-construction benthic community sampling was performed within the proposed north borrow site and reference site for the Dare County shoreline protection project (Versar 2006). Sampling included benthic grabs at ten stations within each of the two sites during the four seasons from spring 2004 through winter 2005. A total of 168 taxa were collected during the study, with polychaetes being the dominant taxa group in occurrence and biomass, especially during the fall and winter. The mean number of taxa present in the fall at the proposed north borrow site was 16 (range of 8 to 26 taxa) and 8 (range of 6 to 11 taxa) in the winter. This is not too different than what was found in the six Topsail Beach borrow sites, where the mean number of species across the six borrow sites ranged from 8 to 17. However, the species richness in the Dare County reference site was significantly greater than the species richness found off Topsail Beach or in the Dare County proposed north borrow site. Mean faunal density in the fall at the Dare County proposed north borrow site (1,555 individuals/m<sup>2</sup>) was slightly higher than the mean density within the Topsail Beach borrow sites (Figure 4, Table 5), while the mean density at the reference site (8,696 individuals/m<sup>2</sup>) was significantly greater. Mean faunal density in the winter was exceptionally higher off Dare County than observed off Topsail Beach, with mean densities of 11,460 individuals/m<sup>2</sup> and 1,398 individuals/m<sup>2</sup> for the reference and borrow sites, respectively. The mean biomass of fall samples was 0.28 g/m<sup>2</sup> and 2.23 g/m<sup>2</sup> for the borrow and reference sites, respectively; as compared to a much higher mean biomass range of 37 g/m<sup>2</sup> to 182 g/m<sup>2</sup> for the six Topsail Beach borrow sites. This comparison also held true for the winter mean biomass values from Dare County.

In summary, the benthic community found within the six proposed borrow sites off Topsail Beach is similar in composition and taxa dominance to those described in other studies along the North Carolina and South Carolina coasts (Byrnes et al. 2003; Versar 2002, 2006; and

Posey and Alphin 2000, 2002). However, the number of species present and abundance were noticeably lower off Topsail Beach than found off Kure Beach (Posey and Alphin 2000) or Dare County (Versar 2006). It is likely that the differences between the benthic community off Topsail Beach, as reported here, and the two referenced studies are due to the more extensive sampling effort associated with baseline monitoring programs, as compared to a less intensive sampling regime for a general characterization study (e.g. ten sampling stations per site off Dare County as compared to three to five stations per site for this study)

## 5.0 LITERATURE CITED

- Byrnes, M.R., R.M. Hammer, B.A. Vittor, S.W. Kelley, D.B. Snyder, J.M. Côté, J.S. Ramsey, T.D. Thibaut, N.W. Phillips, J.D. Wood, and J.D. Germano. 2003. Collection of environmental data within sand resource areas offshore North Carolina and the environmental implications of sand removal for coastal and beach restoration. U.S. Department of the Interior, Minerals Management Service, Leasing Division, Sand and Gravel Unit, Herndon, VA. OCS Report MMS 2000-056, Volume I: Main Text 256 pp. + Volume II: Appendices 69 pp.
- Posey, M.H. and T.D. Alphin. 2000. Monitoring of benthic faunal responses to sediment removal associated with the Carolina Beach and vicinity - Area South Project. Prepared for the Wilmington District, US Army Corps of Engineers, Wilmington, NC. 20 pp. plus appendices.
- Posey, M.H. and T.D. Alphin. 2002. Resilience and stability in an offshore benthic community: Responses to sediment borrow activities and hurricane disturbance. *Journal of Coastal Research*. 18(4): 685-697.
- Versar. 2002. Effects of dredged material beach disposal on surf zone and nearshore fish and benthic resources on Bald Head Island, Caswell Beach, Oak Island, and Holden Beach, North Carolina: Interim study findings. Prepared by Versar, Inc., Columbia, MD, for U.S. Army Corps of Engineers, Wilmington District, Wilmington, NC. 2 volumes.
- Versar. 2006. Year one pre-construction environmental monitoring for the Dare County beach shoreline protection project, Dare County, North Carolina. Prepared by Versar Inc., Columbia, MD, for the U.S. Army Corps of Engineers, Wilmington District, Wilmington, NC. 5 sections plus appendices.

**APPENDIX A**

**Data Quality Control Report**

**Table A-1. Data Quality Control Report****Date:** 13 Nov 06**Wind speed/direction (estimated):** NW @ 10-12 knots**Wave height (estimated):** 1' swells within 2 miles of shore. Beyond 2 miles, 2', steep and building slightly throughout the day.**Water temperature:** 60.9°F**Vessel:** F/V Emma K**Personnel on board:** Capt. Jeff Coward, Duke Hunsaker**Sampling Equipment employed:** Trimble GEO XT handheld gps (sub-meter accuracy)

SI-TEX CVS106 (120 kHz)

Standard Ponar grab

Sample Site	Time	Depth (m)	Sample Coordinates	Sediment Sample	Benthic sample?
F1	10:30	16.3		No	No
F1	10:40	16.3		No	No
F1	11:00	16.3		No	No
F1	11:25	16.3		No	No
F1	11:50	16.3		No	No
B1	12:15	13.4		No	No
B1	12:30	13.4		No	No
B1	12:50	13.4		No	No
B1	1:00	13.4		No	No
B1	1:20	13.4		No	No
B1	1:35	13.4		No	No
B1	1:45	13.4		No	No
B1	2:00	13.4		No	No
B1	2:20	13.4		No	No

Comments: Pot hauler works well in calm water, but sheaves aren't grooved deep enough. When trying to pull the sampler up in rolling sea, the line jumps off the pulley. To avoid injury and/or damage to the vessel, it was necessary to pull the ponar by hand all day. While we have listed 14 sampling attempts, there were numerous drops in between in which misfires occurred or the ponar did not fire. When the ponar did fire properly, it was unable to penetrate the substrate. Only a light dusting of sand was retrieved with each grab. Not enough substrate for analysis. GPS points not taken, since no samples were obtained at the sites. Crossed the bar around 3:00pm. Off water at 3:45pm.

**Table A-1. (continued)****Date:** 28 Nov 06**Wind speed/direction (estimated):** NE @ 10 knots**Wave height (estimated):** 2-3' swells within 2.5 miles of shore. Beyond 2.5 miles, 3-4' rolling swells**Water temperature:** 58.9°F**Vessel:** F/V Emma K**Personnel on board:** Capt. Jeff Coward, Duke Hunsaker, Ryan Lewis**Sampling Equipment employed:** Trimble GEO XT handheld gps (sub-meter accuracy)

SI-TEX CVS106 (120 kHz)

Wildco Shipek grab sampler

Plastic jars/labels prepared for each station

<b>Sample Site</b>	<b>Time</b>	<b>Depth (m)</b>	<b>Sample Coordinates</b>	<b>Sediment Sample</b>	<b>Benthic sample?</b>
A1	8:40	13.7	34°18'16.072" 77°39'47.647"	Yes	Yes
A2	9:05	13.3	34°18'02.486" 77°40'21.868"	Yes	Yes
A3	9:30	13.3	34°18'18.236" 77°39'14.008"	Yes	Yes
A4	9:55	14.2	34°17'54.945" 77°38'53.466"	Yes	Yes
A5	10:30	15.1	34°17'28.093" 77°38'33.163"	Yes	Yes
D1	11:15	14.2	34°18'25.705" 77°36'07.197"	Yes	Yes
D2	11:40	15.3	34°18'13.213" 77°35'43.565"	Yes	Yes
D3	12:00	14.9	34°18'21.953" 77°35'12.038"	Yes	Yes
D4	12:35	14.3	34°18'36.383" 77°35'29.818"	Yes	Yes
E1	1:00	15.7	34°18'54.333" 77°34'14.820"	Yes	Yes
E2	1:25	15.4	34°18'41.374" 77°34'00.078"	Yes	Yes
E3	1:45	14.7	34°18'43.009" 77°33'37.360"	Yes	Yes
E4	2:10	15.3	34°19'01.135" 77°33'54.375"	Yes	Yes
F1	2:40	15.8	34°19'29.216" 77°32'55.069"	Yes	Yes
F2	3:15	15.4	34°19'37.710" 77°32'59.961"	Yes	Yes
F3	3:40	15.5	34°19'44.872" 77°32'40.239"	Yes	Yes
F4	4:15	14.7	34°19'43.171" 77°32'22.118"	Yes	Yes

Comments: Ponar replaced with Wildco Shipek grab sampler. Shipek grab very effective. Sampling went very smoothly. Crossed the bar approximately 5:00pm. Off water at 5:45pm.

**Table A-1. (concluded)****Date:** 29 Nov 06**Wind speed/direction (estimated):** ENE @ 5-10 knots**Wave height (estimated):** 2' swells within 2 miles of shore. Beyond 2 miles, 3' swells with occasional 4' swell. Sea slightly confused.**Water temperature:** 59.2°F**Vessel:** F/V Emma K**Personnel on board:** Capt. Jeff Coward, Duke Hunsaker, Ryan Lewis**Sampling Equipment employed:** Trimble GEO XT handheld gps (sub-meter accuracy)

SI-TEX CVS106 (120 kHz)

Wildco Shipek grab sampler

<b>Sample Site</b>	<b>Time</b>	<b>Depth (m)</b>	<b>Sample Coordinates</b>	<b>Sediment Sample</b>	<b>Benthic sample?</b>
B1	9:30	12.2	34°18'46.266"	Yes	Yes
			77°38'44.909"	Yes	Yes
B2	10:00	13.5	34°18'41.749"	Yes	Yes
			77°38'24.575"	Yes	Yes
B3	10:20	13.2	34°18'57.558"	Yes	Yes
			77°38'37.370"	Yes	Yes
C1	10:55	14.1	34°16'25.432"	Yes	Yes
			77°36'58.081"	Yes	Yes
C2	11:20	14.9	34°16'06.703"	Yes	Yes
			77°37'23.427"	Yes	Yes
C3	11:40	14.6	34°16'18.018"	Yes	Yes
			77°37'44.459"	Yes	Yes
C4	11:55	15.2	34°15'57.406"	Yes	Yes
			77°38'16.160"	Yes	Yes

**Comments:** Sampling went smoothly. No problems. Crossed the bar at approximately 1:00pm. Off water at 1:45pm.

**APPENDIX B**

**Sediment Analysis Report**

=====

TIERRA CONSULTING  
SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

=====

ANALYSIS OF SAMPLE: A-1

PAN=0.52=1.09%  
BVA-DIAL CORDY 2-26-07

SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 47.33

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.54	1.14	98.86	1.14
500.00	1.00	.63	1.33	97.53	2.47
250.00	2.00	4.23	8.94	88.59	11.41
125.00	3.00	35.47	74.94	13.65	86.35
62.50	4.00	5.86	12.38	1.27	98.73

=====

TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE A-1

=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
SAND/SILT CUTOFF 62.50 MICRONS  
SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
PERCENT SAND 98.73

PERCENT SILT AND CLAY 1.27

=====

TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
SHEPARD'S: SAND  
TREFETHEN'S: SAND  
U.S. ARMY CORPS OF ENGINEERS: SAND  
U.S. BUREAU OF SOILS: SAND

=====

PHI PERCENTILES

PHI(05)	1.28 PHI	410.98 MICRONS
PHI(16)	2.06 PHI	239.61 MICRONS
PHI(25)	2.18 PHI	220.47 MICRONS
PHI(50)	2.51 PHI	174.96 MICRONS
PHI(75)	2.85 PHI	138.84 MICRONS
PHI(84)	2.97 PHI	127.75 MICRONS
PHI(95)	3.70 PHI	77.02 MICRONS

---

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*INSUFFICIENT DATA TO PERFORM EXTRAPOLATION\*\*\*

INTERPOLATED PHI PERCENTILES

SAMPLE A-1

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	1.14	98.86
.25	840.90	.8409E+00	1.47	98.53
.50	707.11	.7071E+00	1.81	98.19
.75	594.60	.5946E+00	2.14	97.86
1.00	500.00	.5000E+00	2.47	97.53
1.25	420.45	.4204E+00	4.71	95.29
1.50	353.55	.3536E+00	6.94	93.06
1.75	297.30	.2973E+00	9.17	90.83
2.00	250.00	.2500E+00	11.41	88.59
2.25	210.22	.2102E+00	30.14	69.86
2.50	176.78	.1768E+00	48.88	51.12
2.75	148.65	.1487E+00	67.62	32.38
3.00	125.00	.1250E+00	86.35	13.65
3.25	105.11	.1051E+00	89.45	10.55
3.50	88.39	.8839E-01	92.54	7.46
3.75	74.33	.7433E-01	95.64	4.36
4.00	62.50	.6250E-01	98.73	1.27

STATISTICS AND ENGINEERING PARAMETERS

SAMPLE A-1

INMAN'S STATISTICS:

MEDIAN	2.515 PHI	174.955 MICRONS
MEAN	2.515 PHI	174.955 MICRONS
STANDARD DEVIATION	.454 WELL SORTED	
SKEWNESS	.000 NEAR-SYMMETRICAL	
KURTOSIS	1.662 VERY LEPTOKURTIC	

---

FOLK'S STATISTICS:

MEAN	2.515 PHI	174.955 MICRONS
STANDARD DEVIATION	.593	MODERATELY WELL SORTED
SKEWNESS	-.010	NEAR-SYMMETRICAL
KURTOSIS	1.484	LEPTOKURTIC

---

MOMENT MEASURES:

\*\*INSUFFICIENT DATA FOR MOMENT MEASURES\*\*

---

ENGINEERING PARAMETERS:

D60	2.382 PHI	191.909 MICRONS
D30	2.782 PHI	145.409 MICRONS
D10	3.295 PHI	101.905 MICRONS
COEF. OF UNIFORMITY	1.883	
COEF. OF CURVATURE	1.081	

---

TIERRA CONSULTING  
SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

---

ANALYSIS OF SAMPLE: A-2

PAN=0.26=0.60%  
BVA-DIAL CORDY 2-26-07

SIEVE ANALYSIS

---

SAMPLE WEIGHT = 42.99

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.18	.42	99.58	.42
500.00	1.00	.73	1.70	97.88	2.12
250.00	2.00	5.17	12.03	85.86	14.14
125.00	3.00	31.07	72.27	13.58	86.42
62.50	4.00	5.41	12.58	1.00	99.00

=====  
TEXTURAL CLASSIFICATION AND PHI PERCENTILES                    SAMPLE A-2  
=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF        2000.00 MICRONS  
SAND/SILT CUTOFF          62.50 MICRONS  
SILT/CLAY CUTOFF          4.00 MICRONS

PERCENT GRAVEL            .00  
PERCENT SAND              99.00

PERCENT SILT AND CLAY    1.00

=====  
TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
SHEPARD'S: SAND  
TREFETHEN'S: SAND  
U.S. ARMY CORPS OF ENGINEERS: SAND  
U.S. BUREAU OF SOILS: SAND

=====  
PHI PERCENTILES

PHI(05)	1.24	PHI	423.45	MICRONS
PHI(16)	2.03	PHI	245.59	MICRONS
PHI(25)	2.15	PHI	225.28	MICRONS
PHI(50)	2.50	PHI	177.25	MICRONS
PHI(75)	2.84	PHI	139.46	MICRONS
PHI(84)	2.97	PHI	127.93	MICRONS
PHI(95)	3.68	PHI	77.90	MICRONS

=====

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*INSUFFICIENT DATA TO PERFORM EXTRAPOLATION\*\*\*

=====  
INTERPOLATED PHI PERCENTILES                    SAMPLE A-2  
=====

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	.42	99.58
.25	840.90	.8409E+00	.84	99.16
.50	707.11	.7071E+00	1.27	98.73
.75	594.60	.5946E+00	1.69	98.31
1.00	500.00	.5000E+00	2.12	97.88
1.25	420.45	.4204E+00	5.12	94.88
1.50	353.55	.3536E+00	8.13	91.87
1.75	297.30	.2973E+00	11.14	88.86

2.00	250.00	.2500E+00	14.14	85.86
2.25	210.22	.2102E+00	32.21	67.79
2.50	176.78	.1768E+00	50.28	49.72
2.75	148.65	.1487E+00	68.35	31.65
3.00	125.00	.1250E+00	86.42	13.58
3.25	105.11	.1051E+00	89.56	10.44
3.50	88.39	.8839E-01	92.71	7.29
3.75	74.33	.7433E-01	95.85	4.15
4.00	62.50	.6250E-01	99.00	1.00

## STATISTICS AND ENGINEERING PARAMETERS

## SAMPLE A-2

## INMAN'S STATISTICS:

MEDIAN 2.496 PHI 177.251 MICRONS  
 MEAN 2.496 PHI 177.251 MICRONS  
 STANDARD DEVIATION .470 WELL SORTED  
 SKEWNESS .000 NEAR-SYMMETRICAL  
 KURTOSIS 1.596 VERY LEPTOKURTIC

FOLK'S STATISTICS:

MEAN 2.496 PHI 177.251 MICRONS  
 STANDARD DEVIATION .605 MODERATELY WELL SORTED  
 SKEWNESS -.014 NEAR-SYMMETRICAL  
 KURTOSIS 1.447 LEPTOKURTIC

## MOMENT MEASURES:

\*\* INSUFFICIENT DATA FOR MOMENT MEASURES \*\*

## ENGINEERING PARAMETERS:

D60	2.358	PHI	195.092	MICRONS
D30	2.773	PHI	146.313	MICRONS
D10	3.285	PHI	102.604	MICRONS
COEF. OF UNIFORMITY	1.901			
COEF. OF CURVATURE	1.069			

TIERRA CONSULTING  
IMENTARY PETROLOGY ANALYSIS PROGRAM

**ANALYSIS OF SAMPLE: A-3**

PAN=.30=.6%  
BVA DIAL CORDY 2-26-07

## SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 47.21

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.11	.23	99.77	.23
500.00	1.00	1.15	2.44	97.33	2.67
250.00	2.00	4.71	9.98	87.35	12.65
125.00	3.00	37.05	78.48	8.88	91.12
62.50	4.00	3.82	8.09	.78	99.22

===== TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE A-3 =====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
 SAND/SILT CUTOFF 62.50 MICRONS  
 SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
 PERCENT SAND 99.22

PERCENT SILT AND CLAY .78

## ===== TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
 SHEPARD'S: SAND  
 TREFETHEN'S: SAND  
 U.S. ARMY CORPS OF ENGINEERS: SAND  
 U.S. BUREAU OF SOILS: SAND

## ===== PHI PERCENTILES

PHI(05)	1.23	PHI	425.24	MICRONS
PHI(16)	2.04	PHI	242.70	MICRONS
PHI(25)	2.16	PHI	224.16	MICRONS
PHI(50)	2.48	PHI	179.74	MICRONS
PHI(75)	2.79	PHI	144.13	MICRONS
PHI(84)	2.91	PHI	133.12	MICRONS
PHI(95)	3.48	PHI	89.69	MICRONS

===== EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*INSUFFICIENT DATA TO PERFORM EXTRAPOLATION\*\*\*

## INTERPOLATED PHI PERCENTILES

SAMPLE A-3

PHI	MICRONS	MM	PCT.	RET	PCT. FINER
.00	1000.00	.1000E+01	.23		99.77
.25	840.90	.8409E+00	.84		99.16
.50	707.11	.7071E+00	1.45		98.55
.75	594.60	.5946E+00	2.06		97.94
1.00	500.00	.5000E+00	2.67		97.33
1.25	420.45	.4204E+00	5.16		94.84
1.50	353.55	.3536E+00	7.66		92.34
1.75	297.30	.2973E+00	10.15		89.85
2.00	250.00	.2500E+00	12.65		87.35
2.25	210.22	.2102E+00	32.27		67.73
2.50	176.78	.1768E+00	51.89		48.11
2.75	148.65	.1487E+00	71.50		28.50
3.00	125.00	.1250E+00	91.12		8.88
3.25	105.11	.1051E+00	93.15		6.85
3.50	88.39	.8839E-01	95.17		4.83
3.75	74.33	.7433E-01	97.19		2.81
4.00	62.50	.6250E-01	99.22		.78

## STATISTICS AND ENGINEERING PARAMETERS

SAMPLE A-3

TINMAN'S STATISTICS:

MEDIAN 2.476 PHI 179.745 MICRONS  
 MEAN 2.476 PHI 179.745 MICRONS  
 STANDARD DEVIATION .433 WELL SORTED  
 SKEWNESS .000 NEAR-SYMMETRICAL  
 KURTOSIS 1.591 VERY LEPTOKURTIC

FOLK'S STATISTICS:

MEAN 2.476 PHI 179.745 MICRONS  
 STANDARD DEVIATION .557 MODERATELY WELL SORTED  
 SKEWNESS -.053 NEAR-SYMMETRICAL  
 KURTOSIS 1.444 LEPTOKURTIC

#### MOMENT MEASURES:

\*\*INSUFFICIENT DATA FOR MOMENT MEASURES\*\*

#### ENGINEERING PARAMETERS:

D60	2.349	PHI	196.342	MICRONS
D30	2.731	PHI	150.640	MICRONS
D10	2.986	PHT	126.248	MICRONS

COEF. OF UNIFORMITY 1.555  
COEF. OF CURVATURE .915

=====  
TIERRA CONSULTING  
SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM  
=====

ANALYSIS OF SAMPLE: A-4

PAN=0.07=.16%  
BVA-DIAL CORDY 2-26-07

SIEVE ANALYSIS

SAMPLE WEIGHT = 42.79

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.30	.70	99.30	.70
500.00	1.00	1.52	3.55	95.75	4.25
250.00	2.00	5.33	12.46	83.29	16.71
125.00	3.00	30.28	70.76	12.53	87.47
62.50	4.00	5.24	12.25	.28	99.72

=====  
TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE A-4  
=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
SAND/SILT CUTOFF 62.50 MICRONS  
SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
PERCENT SAND 99.72

PERCENT SILT AND CLAY .28

=====  
TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
SHEPARD'S: SAND  
TREFETHEN'S: SAND  
U.S. ARMY CORPS OF ENGINEERS: SAND  
U.S. BUREAU OF SOILS: SAND

## PHI PERCENTILES

PHI(05)	1.06	PHI	479.65	MICRONS
PHI(16)	1.94	PHI	260.07	MICRONS
PHI(25)	2.12	PHI	230.50	MICRONS
PHI(50)	2.47	PHI	180.44	MICRONS
PHI(75)	2.82	PHI	141.24	MICRONS
PHI(84)	2.95	PHI	129.33	MICRONS
PHI(95)	3.61	PHI	81.64	MICRONS

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

## INTERPOLATED PHI PERCENTILES

SAMPLE A-4

PHI	MICRONS	MM	PCT.	RET	PCT. FINER
.00	1000.00	.1000E+01	.70		99.30
.25	840.90	.8409E+00	1.59		98.41
.50	707.11	.7071E+00	2.48		97.52
.75	594.60	.5946E+00	3.37		96.63
1.00	500.00	.5000E+00	4.25		95.75
1.25	420.45	.4204E+00	7.37		92.63
1.50	353.55	.3536E+00	10.48		89.52
1.75	297.30	.2973E+00	13.60		86.40
2.00	250.00	.2500E+00	16.71		83.29
2.25	210.22	.2102E+00	34.40		65.60
2.50	176.78	.1768E+00	52.09		47.91
2.75	148.65	.1487E+00	69.78		30.22
3.00	125.00	.1250E+00	87.47		12.53
3.25	105.11	.1051E+00	90.54		9.46
3.50	88.39	.8839E-01	93.60		6.40
3.75	74.33	.7433E-01	96.66		3.34
4.00	62.50	.6250E-01	99.72		.28

STATISTICS AND ENGINEERING PARAMETERS

SAMPLE A-4

## TINMAN'S STATISTICS:

MEDIAN	2.470	PHI	180.436	MICRONS
MEAN	2.447	PHI	183.395	MICRONS
STANDARD DEVIATION	.504	MODERATELY WELL SORTED		
SKEWNESS	-.047	NEAR-SYMMETRICAL		
KURTOSIS	1.535	VERY LEPTOKURTIC		

FOLK'S STATISTICS:

MEAN 2.455 PHI 182.403 MICRONS  
 STANDARD DEVIATION .639 MODERATELY WELL SORTED  
 SKEWNESS -.075 NEAR-SYMMETRICAL  
 KURTOSIS 1.482 LEPTOKURTIC

=====

MOMENT MEASURES:

MEAN	2.399 PHI	189.621 MICRONS
SECOND MOMENT	.443	
STANDARD DEVIATION	.666	MODERATELY WELL SORTED
THIRD MOMENT	-.340	
SKEWNESS	-1.152	STRONGLY COARSE-SKewed
FOURTH MOMENT	1.218	
KURTOSIS	2.066	VERY LEPTOKURTIC

=====

ENGINEERING PARAMETERS:

D60	2.329 PHI	199.004 MICRONS
D30	2.753 PHI	148.335 MICRONS
D10	3.206 PHI	108.345 MICRONS
COEF. OF UNIFORMITY	1.837	
COEF. OF CURVATURE	1.021	

=====

TIERRA CONSULTING  
SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

=====

ANALYSIS OF SAMPLE: A-5

PAN=0.05=.11%

BVA DIAL CORDY 2-26-07

SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 45.18

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.17	.38	99.62	.38
500.00	1.00	.49	1.08	98.54	1.46
250.00	2.00	7.30	16.16	82.38	17.62
125.00	3.00	33.57	74.30	8.08	91.92
62.50	4.00	3.60	7.97	.11	99.89

=====  
TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE A-5  
=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
SAND/SILT CUTOFF 62.50 MICRONS  
SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
PERCENT SAND 99.89

PERCENT SILT AND CLAY .11

=====

TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
SHEPARD'S: SAND  
TREFETHEN'S: SAND  
U.S. ARMY CORPS OF ENGINEERS: SAND  
U.S. BUREAU OF SOILS: SAND

=====

PHI PERCENTILES

PHI(05)	1.22	PHI	429.57	MICRONS
PHI(16)	1.90	PHI	267.97	MICRONS
PHI(25)	2.10	PHI	233.36	MICRONS
PHI(50)	2.44	PHI	184.82	MICRONS
PHI(75)	2.77	PHI	146.37	MICRONS
PHI(84)	2.89	PHI	134.59	MICRONS
PHI(95)	3.39	PHI	95.63	MICRONS

=====

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

=====  
INTERPOLATED PHI PERCENTILES SAMPLE A-5  
=====

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	.38	99.62
.25	840.90	.8409E+00	.65	99.35
.50	707.11	.7071E+00	.92	99.08
.75	594.60	.5946E+00	1.19	98.81
1.00	500.00	.5000E+00	1.46	98.54
1.25	420.45	.4204E+00	5.50	94.50
1.50	353.55	.3536E+00	9.54	90.46
1.75	297.30	.2973E+00	13.58	86.42
2.00	250.00	.2500E+00	17.62	82.38

2.25	210.22	.2102E+00	36.19	63.81
2.50	176.78	.1768E+00	54.77	45.23
2.75	148.65	.1487E+00	73.35	26.65
3.00	125.00	.1250E+00	91.92	8.08
3.25	105.11	.1051E+00	93.91	6.09
3.50	88.39	.8839E-01	95.91	4.09
3.75	74.33	.7433E-01	97.90	2.10
4.00	62.50	.6250E-01	99.89	.11

===== STATISTICS AND ENGINEERING PARAMETERS =====

SAMPLE A-5

INMAN'S STATISTICS:

MEDIAN	2.436	PHI	184.820	MICRONS
MEAN	2.397	PHI	189.910	MICRONS
STANDARD DEVIATION	.497	WELL SORTED		
SKEWNESS	-.079	NEAR-SYMMETRICAL		
KURTOSIS	1.181	LEPTOKURTIC		

FOLK'S STATISTICS:

MEAN	2.410	PHI	188.198	MICRONS
STANDARD DEVIATION	.577	MODERATELY WELL SORTED		
SKEWNESS	-.101	COARSE-SKewed		
KURTOSIS	1.320	LEPTOKURTIC		

MOMENT MEASURES:

MEAN	2.382	PHI	191.795	MICRONS
SECOND MOMENT	.305			
STANDARD DEVIATION	.553	MODERATELY WELL SORTED		
THIRD MOMENT	-.161			
SKEWNESS	-.954	STRONGLY COARSE-SKewed		
FOURTH MOMENT	.618			
KURTOSIS	2.211	VERY LEPTOKURTIC		

ENGINEERING PARAMETERS:

D60	2.301	PHI	202.891	MICRONS
D30	2.705	PHI	153.363	MICRONS
D10	2.974	PHI	127.260	MICRONS
COEF. OF UNIFORMITY	1.594			
COEF. OF CURVATURE	.911			

=====  
 TIERRA CONSULTING  
 SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM  
 =====

TIERRA CONSULTING  
SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

---

ANALYSIS OF SAMPLE: B-1

PAN=0.00  
BVA DIAL CORDY 2-26-07

SIEVE ANALYSIS

---

SAMPLE WEIGHT = 48.25

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	24.92	51.65	48.35	51.65
500.00	1.00	8.30	17.20	31.15	68.85
250.00	2.00	9.19	19.05	12.10	87.90
125.00	3.00	5.56	11.52	.58	99.42
62.50	4.00	.43	.89	-.31	100.31

\*\*\*\*\*  
\* \*\*\*ERROR\*\*\* \*\*\*ERROR\*\*\* CUMULATIVE PERCENT RETAINED GREATER THAN 100% \*  
\*  
\*\*\*\*\*

=====  
TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE B-1  
=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
SAND/SILT CUTOFF 62.50 MICRONS  
SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
PERCENT SAND 100.31

PERCENT SILT AND CLAY -.31

TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
SHEPARD'S: SAND  
TREFETHEN'S: SAND  
U.S. ARMY CORPS OF ENGINEERS: SAND  
U.S. BUREAU OF SOILS: SAND

---

PHI PERCENTILES

PHI(75)	1.32 PHI	399.73 MICRONS
PHI(84)	1.80 PHI	288.09 MICRONS
PHI(95)	2.62 PHI	163.07 MICRONS

---

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

INTERPOLATED PHI PERCENTILES SAMPLE B-1

---

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	51.65	48.35
.25	840.90	.8409E+00	55.95	44.05
.50	707.11	.7071E+00	60.25	39.75
.75	594.60	.5946E+00	64.55	35.45
1.00	500.00	.5000E+00	68.85	31.15
1.25	420.45	.4204E+00	73.61	26.39
1.50	353.55	.3536E+00	78.37	21.63
1.75	297.30	.2973E+00	83.13	16.87
2.00	250.00	.2500E+00	87.90	12.10
2.25	210.22	.2102E+00	90.78	9.22
2.50	176.78	.1768E+00	93.66	6.34
2.75	148.65	.1487E+00	96.54	3.46
3.00	125.00	.1250E+00	99.42	.58
3.25	105.11	.1051E+00	99.64	.36
3.50	88.39	.8839E-01	99.87	.13
3.75	74.33	.7433E-01	100.09	-.09
4.00	62.50	.6250E-01	100.31	-.31

---

STATISTICS AND ENGINEERING PARAMETERS SAMPLE B-1

---

\*\*INSUFFICIENT DATA FOR ANY STATISTICAL CALCULATIONS\*\*

---

ENGINEERING PARAMETERS:

D60 IS NOT CALCULATED--FIRST SIEVE RETAINS OVER 40%  
 D30 1.060 PHI 479.502 MICRONS  
 D10 2.183 PHI 220.285 MICRONS  
 COEF. OF UNIFORMITY IS NOT CALCULATED  
 COEF. OF CURVATURE IS NOT CALCULATED

TIERRA CONSULTING  
SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

## ANALYSIS OF SAMPLE: B-2

PAN=.05=.10%  
BVA-DIAL CORDY 2-26-07

## SIEVE ANALYSIS

SAMPLE WEIGHT = 49.36

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.55	1.11	98.89	1.11
500.00	1.00	1.62	3.28	95.60	4.40
250.00	2.00	7.73	15.66	79.94	20.06
125.00	3.00	32.93	66.71	13.23	86.77
62.50	4.00	3.91	7.92	5.31	94.69

## TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE B-2

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF        2000.00 MICRONS  
 SAND/SILT CUTOFF        62.50 MICRONS  
 SILT/CLAY CUTOFF        4.00 MICRONS

PERCENT GRAVEL	.00
PERCENT SAND	94.69
PERCENT SILT AND CLAY	5.31

## TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
SHEPARD'S: SAND  
TREFETHEN'S: SAND  
U.S. ARMY CORPS OF ENGINEERS: SAND  
U.S. BUREAU OF SOILS: SAND

PHI PERCENTILES

PHI(05)	1.04 PHI	486.82 MICRONS
PHI(16)	1.74 PHI	299.17 MICRONS
PHI(25)	2.07 PHI	237.48 MICRONS
PHI(50)	2.45 PHI	183.16 MICRONS
PHI(75)	2.82 PHI	141.26 MICRONS
PHI(84)	2.96 PHI	128.65 MICRONS

---

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*INSUFFICIENT DATA TO PERFORM EXTRAPOLATION\*\*\*

INTERPOLATED PHI PERCENTILES

SAMPLE B-2

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	1.11	98.89
.25	840.90	.8409E+00	1.93	98.07
.50	707.11	.7071E+00	2.76	97.24
.75	594.60	.5946E+00	3.58	96.42
1.00	500.00	.5000E+00	4.40	95.60
1.25	420.45	.4204E+00	8.31	91.69
1.50	353.55	.3536E+00	12.23	87.77
1.75	297.30	.2973E+00	16.14	83.86
2.00	250.00	.2500E+00	20.06	79.94
2.25	210.22	.2102E+00	36.74	63.26
2.50	176.78	.1768E+00	53.41	46.59
2.75	148.65	.1487E+00	70.09	29.91
3.00	125.00	.1250E+00	86.77	13.23
3.25	105.11	.1051E+00	88.75	11.25
3.50	88.39	.8839E-01	90.73	9.27
3.75	74.33	.7433E-01	92.71	7.29
4.00	62.50	.6250E-01	94.69	5.31

STATISTICS AND ENGINEERING PARAMETERS

SAMPLE B-2

INMAN'S STATISTICS:

MEDIAN	2.449 PHI	183.159 MICRONS
MEAN	2.350 PHI	196.185 MICRONS
STANDARD DEVIATION	.609	MODERATELY WELL SORTED
SKEWNESS	-.163	COARSE-SKewed

---

FOLK'S STATISTICS:

MEAN	2.383 PHI	191.743 MICRONS
------	-----------	-----------------

---

MOMENT MEASURES:

\*\*INSUFFICIENT DATA FOR MOMENT MEASURES\*\*

---

ENGINEERING PARAMETERS:

D60	2.299 PHI	203.213 MICRONS
D30	2.749 PHI	148.793 MICRONS
D10	3.408 PHI	94.230 MICRONS
COEF. OF UNIFORMITY	2.157	
COEF. OF CURVATURE	1.156	

---

TIERRA CONSULTING

SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

---

ANALYSIS OF SAMPLE: B-3

PAN=0.11=0.25%

BVA DIAL CORDY 2-26-07

SIEVE ANALYSIS

---

SAMPLE WEIGHT = 43.90

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	27.79	63.30	36.70	63.30
500.00	1.00	9.11	20.75	15.95	84.05
250.00	2.00	3.80	8.66	7.29	92.71
125.00	3.00	2.70	6.15	1.14	98.86
62.50	4.00	.32	.73	.41	99.59

---

TEXTURAL CLASSIFICATION AND PHI PERCENTILES                   SAMPLE B-3

---

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF	2000.00 MICRONS
SAND/SILT CUTOFF	62.50 MICRONS
SILT/CLAY CUTOFF	4.00 MICRONS
PERCENT GRAVEL	.00
PERCENT SAND	99.59

PERCENT SILT AND CLAY .41

---

## TEXTURAL DESCRIPTIONS

FOLK'S: SAND

SHEPARD'S: SAND

TREFETHEN'S: SAND

U.S. ARMY CORPS OF ENGINEERS: SAND

U. S. BUREAU OF STANDARDS

## PHI PERCENTILES

PHI(75)	.56	PHI	676.58	MICRONS
PHI(84)	1.00	PHI	500.91	MICRONS
PHI(95)	2.37	PHI	193.15	MICRONS

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

## INTERPOLATED PHI PERCENTILES

SAMPLE B-3

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	63.30	36.70
.25	840.90	.8409E+00	68.49	31.51
.50	707.11	.7071E+00	73.68	26.32
.75	594.60	.5946E+00	78.87	21.13
1.00	500.00	.5000E+00	84.05	15.95
1.25	420.45	.4204E+00	86.22	13.78
1.50	353.55	.3536E+00	88.38	11.62
1.75	297.30	.2973E+00	90.55	9.45
2.00	250.00	.2500E+00	92.71	7.29
2.25	210.22	.2102E+00	94.25	5.75
2.50	176.78	.1768E+00	95.79	4.21
2.75	148.65	.1487E+00	97.32	2.68
3.00	125.00	.1250E+00	98.86	1.14
3.25	105.11	.1051E+00	99.04	.96
3.50	88.39	.8839E-01	99.23	.77
3.75	74.33	.7433E-01	99.41	.59
4.00	62.50	.6250E-01	99.59	.41

## STATISTICS AND ENGINEERING PARAMETERS

SAMPLE B-3

\*\* INSUFFICIENT DATA FOR ANY STATISTICAL CALCULATIONS \*\*

## ENGINEERING PARAMETERS:

P60 IS NOT CALCULATED--FIRST SIEVE RETAINS OVER 40%

D30 .323 PHI 799.560 MICRONS

D10 1.687 PHI

COEFF. OF UNIFORMITY IS NOT CALCULATED

COEF. OF UNIFORMITY IS NOT CALCULATED  
COEF. OF CURVATURE IS NOT CALCULATED

=====

TIERRA CONSULTING  
SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

=====

ANALYSIS OF SAMPLE: C-3

PAN=.08=.17%  
BVA- DIAL CORDY 2-26-07

SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 44.59

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.02	.04	99.96	.04
500.00	1.00	.28	.63	99.33	.67
250.00	2.00	5.58	12.51	86.81	13.19
125.00	3.00	35.30	79.17	7.65	92.35
62.50	4.00	3.31	7.42	.22	99.78

=====

TEXTURAL CLASSIFICATION AND PHI PERCENTILES                    SAMPLE C-3

=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF        2000.00 MICRONS  
SAND/SILT CUTOFF           62.50 MICRONS  
SILT/CLAY CUTOFF           4.00 MICRONS

PERCENT GRAVEL             .00  
PERCENT SAND              99.78

PERCENT SILT AND CLAY     .22

=====

TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
SHEPARD'S: SAND  
TREFETHEN'S: SAND  
U.S. ARMY CORPS OF ENGINEERS: SAND  
U.S. BUREAU OF SOILS: SAND

=====

PHI PERCENTILES

PHI(05)	1.35 PHI	393.44 MICRONS
PHI(16)	2.04 PHI	243.92 MICRONS
PHI(25)	2.15 PHI	225.43 MICRONS
PHI(50)	2.47 PHI	181.12 MICRONS
PHI(75)	2.78 PHI	145.51 MICRONS
PHI(84)	2.89 PHI	134.48 MICRONS
PHI(95)	3.36 PHI	97.62 MICRONS

---

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

INTERPOLATED PHI PERCENTILES

SAMPLE C-3

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	.04	99.96
.25	840.90	.8409E+00	.20	99.80
.50	707.11	.7071E+00	.36	99.64
.75	594.60	.5946E+00	.52	99.48
1.00	500.00	.5000E+00	.67	99.33
1.25	420.45	.4204E+00	3.80	96.20
1.50	353.55	.3536E+00	6.93	93.07
1.75	297.30	.2973E+00	10.06	89.94
2.00	250.00	.2500E+00	13.19	86.81
2.25	210.22	.2102E+00	32.98	67.02
2.50	176.78	.1768E+00	52.77	47.23
2.75	148.65	.1487E+00	72.56	27.44
3.00	125.00	.1250E+00	92.35	7.65
3.25	105.11	.1051E+00	94.21	5.79
3.50	88.39	.8839E-01	96.06	3.94
3.75	74.33	.7433E-01	97.92	2.08
4.00	62.50	.6250E-01	99.78	.22

STATISTICS AND ENGINEERING PARAMETERS

SAMPLE C-3

INMAN'S STATISTICS:

MEDIAN	2.465 PHI	181.116 MICRONS
MEAN	2.465 PHI	181.116 MICRONS
STANDARD DEVIATION	.429 WELL SORTED	
SKEWNESS	.000 NEAR-SYMMETRICAL	
KURTOSIS	1.341 LEPTOKURTIC	

---

FOLK'S STATISTICS:

MEAN	2.465 PHI	181.116 MICRONS
STANDARD DEVIATION	.519	MODERATELY WELL SORTED
SKEWNESS	-.057	NEAR-SYMMETRICAL
KURTOSIS	1.305	LEPTOKURTIC

---

MOMENT MEASURES:

MEAN	2.430 PHI	185.619 MICRONS
SECOND MOMENT	.224	
STANDARD DEVIATION	.474	WELL SORTED
THIRD MOMENT	-.066	
SKEWNESS	-.617	STRONGLY COARSE-SKewed
FOURTH MOMENT	.311	
KURTOSIS	2.060	VERY LEPTOKURTIC

---

ENGINEERING PARAMETERS:

D60	2.339 PHI	197.689 MICRONS
D30	2.718 PHI	152.022 MICRONS
D10	2.970 PHI	127.601 MICRONS
COEF. OF UNIFORMITY	1.549	
COEF. OF CURVATURE	.916	

---

TIERRA CONSULTING

SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

---

ANALYSIS OF SAMPLE: C-4

PAN=.06=.12%

BVA- DIAL CORDY 2-26-07

SIEVE ANALYSIS

---

SAMPLE WEIGHT = 48.86

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.06	.12	99.88	.12
500.00	1.00	1.00	2.05	97.83	2.17
250.00	2.00	4.59	9.39	88.44	11.56
125.00	3.00	40.06	81.99	6.45	93.55
62.50	4.00	2.94	6.02	.43	99.57

=====  
TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE C-4  
=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
SAND/SILT CUTOFF 62.50 MICRONS  
SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
PERCENT SAND 99.57  
PERCENT SILT AND CLAY .43

=====

TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
SHEPARD'S: SAND  
TREFETHEN'S: SAND  
U.S. ARMY CORPS OF ENGINEERS: SAND  
U.S. BUREAU OF SOILS: SAND

=====

PHI PERCENTILES

PHI(05)	1.30 PHI	405.76 MICRONS
PHI(16)	2.05 PHI	240.80 MICRONS
PHI(25)	2.16 PHI	223.16 MICRONS
PHI(50)	2.47 PHI	180.64 MICRONS
PHI(75)	2.77 PHI	146.23 MICRONS
PHI(84)	2.88 PHI	135.51 MICRONS
PHI(95)	3.24 PHI	105.81 MICRONS

=====

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

=====  
INTERPOLATED PHI PERCENTILES SAMPLE C-4  
=====

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	.12	99.88
.25	840.90	.8409E+00	.63	99.37
.50	707.11	.7071E+00	1.15	98.85
.75	594.60	.5946E+00	1.66	98.34
1.00	500.00	.5000E+00	2.17	97.83
1.25	420.45	.4204E+00	4.52	95.48
1.50	353.55	.3536E+00	6.87	93.13
1.75	297.30	.2973E+00	9.22	90.78
2.00	250.00	.2500E+00	11.56	88.44
2.25	210.22	.2102E+00	32.06	67.94
2.50	176.78	.1768E+00	52.56	47.44

2.75	148.65	.1487E+00	73.06	26.94
3.00	125.00	.1250E+00	93.55	6.45
3.25	105.11	.1051E+00	95.06	4.94
3.50	88.39	.8839E-01	96.56	3.44
3.75	74.33	.7433E-01	98.07	1.93
4.00	62.50	.6250E-01	99.57	.43

## STATISTICS AND ENGINEERING PARAMETERS

SAMPLE C-4

## INMAN'S STATISTICS:

MEDIAN 2.469 PHI 180.642 MICRONS  
 MEAN 2.469 PHI 180.642 MICRONS  
 STANDARD DEVIATION .415 WELL SORTED  
 SKEWNESS .000 NEAR-SYMMETRICAL  
 KURTOSIS 1.338 LEPTOKURTIC

## FOLK'S STATISTICS:

MEAN 2.469 PHI 180.642 MICRONS  
STANDARD DEVIATION .501 MODERATELY WELL SORTED  
SKEWNESS -.102 COARSE-SKewed  
KURTOSIS 1.303 LEPTOKURTIC

## MOMENT MEASURES:

MEAN	2.411	PHI	188.043	MICRONS
SECOND MOMENT	.241			
STANDARD DEVIATION	.491	WELL SORTED		
THIRD MOMENT	-.166			
SKEWNESS	-1.401	STRONGLY COARSE-SKewed		
FOURTH MOMENT	.510			
KURTOSIS	2.930	VERY LEPTOKURTIC		

## ENGINEERING PARAMETERS:

D60	2.347	PHI	196.578	MICRONS
D30	2.713	PHI	152.541	MICRONS
D10	2.957	PHI	128.812	MICRONS
COEF. OF UNIFORMITY	1.526			
COEF. OF CURVATURE	919			

TIERRA CONSULTING

SEDPET LIT: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

ANALYSIS OF SAMPLE: C1

$$PAN = .03 = .06\%$$

BVA DIAL CORDY 2-26-07

## SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 48.84

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
500.00	1.00	1.12	2.29	97.71	2.29
250.00	2.00	7.65	15.66	82.04	17.96
125.00	3.00	37.92	77.64	4.40	95.60
62.50	4.00	2.11	4.32	.08	99.92

=====  
TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE C1  
=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
 SAND/SILT CUTOFF 62.50 MICRONS  
 SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
 PERCENT SAND 99.92

PERCENT SILT AND CLAY .08

=====  
TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
 SHEPARD'S: SAND  
 TREFETHEN'S: SAND  
 U.S. ARMY CORPS OF ENGINEERS: SAND  
 U.S. BUREAU OF SOILS: SAND

=====  
PHI PERCENTILES

PHI(05)	1.17 PHI	443.56 MICRONS
PHI(16)	1.88 PHI	272.61 MICRONS
PHI(25)	2.09 PHI	234.76 MICRONS
PHI(50)	2.41 PHI	187.80 MICRONS
PHI(75)	2.73 PHI	150.24 MICRONS
PHI(84)	2.85 PHI	138.64 MICRONS
PHI(95)	2.99 PHI	125.67 MICRONS

=====  
EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

## INTERPOLATED PHI PERCENTILES

SAMPLE C1

PHI	MICRONS	MM	PCT. RET	PCT. FINER
1.00	500.00	.5000E+00	2.29	97.71
1.25	420.45	.4204E+00	6.21	93.79
1.50	353.55	.3536E+00	10.12	89.88
1.75	297.30	.2973E+00	14.04	85.96
2.00	250.00	.2500E+00	17.96	82.04
2.25	210.22	.2102E+00	37.37	62.63
2.50	176.78	.1768E+00	56.78	43.22
2.75	148.65	.1487E+00	76.19	23.81
3.00	125.00	.1250E+00	95.60	4.40
3.25	105.11	.1051E+00	96.68	3.32
3.50	88.39	.8839E-01	97.76	2.24
3.75	74.33	.7433E-01	98.84	1.16
4.00	62.50	.6250E-01	99.92	.08

## STATISTICS AND ENGINEERING PARAMETERS

SAMPLE C1

## INMAN'S STATISTICS:

MEDIAN	2.413	PHI	187.803	MICRONS
MEAN	2.363	PHI	194.406	MICRONS
STANDARD DEVIATION	.488	WELL SORTED		
SKEWNESS	-.102	COARSE-SKewed		
KURTOSIS	.865	PLATYKURTIC		

FOLK'S STATISTICS:

MEAN 2.379 PHI 192.180 MICRONS  
STANDARD DEVIATION .520 MODERATELY WELL SORTED  
SKEWNESS -.233 COARSE-SKewed  
KURTOSIS 1.158 LEPTOKURTIC

#### MOMENT MEASURES:

MEAN	2.339	PHI	197.694	MICRONS
SECOND MOMENT	.266			
STANDARD DEVIATION	.516	MODERATELY WELL SORTED		
THIRD MOMENT	-.164			
SKEWNESS	-1.194	STRONGLY COARSE-SKewed		
FOURTH MOMENT	.419			
KURTOSIS	1.970	VERY LEPTOKURTIC		

ENGINEERING PARAMETERS:

D60	2.284 PHI	205.340 MICRONS
D30	2.670 PHI	157.093 MICRONS
D10	2.928 PHI	131.406 MICRONS
COEF. OF UNIFORMITY	1.563	
COEF. OF CURVATURE	.915	

TIERRA CONSULTING

SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

=====

ANALYSIS OF SAMPLE: C-2

PAN=0.06g=.12%

BVA DIAL CORDY 2-26-07

SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 49.58

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.07	.14	99.86	.14
500.00	1.00	.75	1.51	98.35	1.65
250.00	2.00	6.78	13.67	84.67	15.33
125.00	3.00	38.64	77.93	6.74	93.26
62.50	4.00	3.25	6.56	.18	99.82

=====

TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE C-2

=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF	2000.00 MICRONS
SAND/SILT CUTOFF	62.50 MICRONS
SILT/CLAY CUTOFF	4.00 MICRONS

PERCENT GRAVEL	.00
PERCENT SAND	99.82

PERCENT SILT AND CLAY	.18
-----------------------	-----

TEXTURAL DESCRIPTIONS

FOLK'S: SAND

SHEPARD'S: SAND

TREFETHEN'S: SAND

U.S. ARMY CORPS OF ENGINEERS: SAND

U.S. BUREAU OF SOILS: SAND

=====

PHI PERCENTILES

PHI(05)	1.24	PHI	422.00	MICRONS
PHI(16)	2.01	PHI	248.51	MICRONS
PHI(25)	2.12	PHI	229.39	MICRONS
PHI(50)	2.44	PHI	183.66	MICRONS
PHI(75)	2.77	PHI	147.05	MICRONS
PHI(84)	2.88	PHI	135.73	MICRONS
PHI(95)	3.26	PHI	104.03	MICRONS

=====

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

=====

INTERPOLATED PHI PERCENTILES

SAMPLE C-2

=====

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	.14	99.86
.25	840.90	.8409E+00	.52	99.48
.50	707.11	.7071E+00	.90	99.10
.75	594.60	.5946E+00	1.28	98.72
1.00	500.00	.5000E+00	1.65	98.35
1.25	420.45	.4204E+00	5.07	94.93
1.50	353.55	.3536E+00	8.49	91.51
1.75	297.30	.2973E+00	11.91	88.09
2.00	250.00	.2500E+00	15.33	84.67
2.25	210.22	.2102E+00	34.81	65.19
2.50	176.78	.1768E+00	54.30	45.70
2.75	148.65	.1487E+00	73.78	26.22
3.00	125.00	.1250E+00	93.26	6.74
3.25	105.11	.1051E+00	94.90	5.10
3.50	88.39	.8839E-01	96.54	3.46
3.75	74.33	.7433E-01	98.18	1.82
4.00	62.50	.6250E-01	99.82	.18

=====

STATISTICS AND ENGINEERING PARAMETERS

SAMPLE C-2

=====

INMAN'S STATISTICS:

MEDIAN	2.445 PHI	183.662 MICRONS
MEAN	2.445 PHI	183.662 MICRONS
STANDARD DEVIATION	.436	WELL SORTED
SKEWNESS	.000	NEAR-SYMMETRICAL
KURTOSIS	1.315	LEPTOKURTIC

=====

FOLK'S STATISTICS:

MEAN	2.445 PHI	183.662 MICRONS
STANDARD DEVIATION	.524	MODERATELY WELL SORTED
SKEWNESS	-.094	NEAR-SYMMETRICAL
KURTOSIS	1.291	LEPTOKURTIC

=====

MOMENT MEASURES:

MEAN	2.390 PHI	190.812 MICRONS
SECOND MOMENT	.264	
STANDARD DEVIATION	.514	MODERATELY WELL SORTED
THIRD MOMENT	-.142	
SKEWNESS	-1.043	STRONGLY COARSE-SKewed
FOURTH MOMENT	.477	
KURTOSIS	2.274	VERY LEPTOKURTIC

=====

ENGINEERING PARAMETERS:

D60	2.317 PHI	200.745 MICRONS
D30	2.702 PHI	153.733 MICRONS
D10	2.958 PHI	128.681 MICRONS
COEF. OF UNIFORMITY	1.560	
COEF. OF CURVATURE	.915	

=====

TIERRA CONSULTING  
SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

=====

ANALYSIS OF SAMPLE: D-1

PAN=0.05=<5%  
BVA DIAL CORDY 2-26-07

## SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 49.48

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.21	.42	99.58	.42
500.00	1.00	2.48	5.01	94.56	5.44
250.00	2.00	13.61	27.51	67.06	32.94
125.00	3.00	31.55	63.76	3.29	96.71
62.50	4.00	1.56	3.15	.14	99.86

===== TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE D-1 =====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
 SAND/SILT CUTOFF 62.50 MICRONS  
 SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
 PERCENT SAND 99.86

PERCENT SILT AND CLAY .14

## TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
 SHEPARD'S: SAND  
 TREFETHEN'S: SAND  
 U.S. ARMY CORPS OF ENGINEERS: SAND  
 U.S. BUREAU OF SOILS: SAND

## PHI PERCENTILES

PHI(05)	.91 PHI	531.12 MICRONS
PHI(16)	1.38 PHI	383.14 MICRONS
PHI(25)	1.71 PHI	305.40 MICRONS
PHI(50)	2.27 PHI	207.69 MICRONS
PHI(75)	2.66 PHI	158.26 MICRONS
PHI(84)	2.80 PHI	143.51 MICRONS
PHI(95)	2.97 PHI	127.34 MICRONS

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

## INTERPOLATED PHT PERCENTILES

SAMPLE D-1

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	.42	99.58
.25	840.90	.8409E+00	1.68	98.32
.50	707.11	.7071E+00	2.93	97.07
.75	594.60	.5946E+00	4.18	95.82
1.00	500.00	.5000E+00	5.44	94.56
1.25	420.45	.4204E+00	12.31	87.69
1.50	353.55	.3536E+00	19.19	80.81
1.75	297.30	.2973E+00	26.07	73.93
2.00	250.00	.2500E+00	32.94	67.06
2.25	210.22	.2102E+00	48.88	51.12
2.50	176.78	.1768E+00	64.82	35.18
2.75	148.65	.1487E+00	80.76	19.24
3.00	125.00	.1250E+00	96.71	3.29
3.25	105.11	.1051E+00	97.49	2.51
3.50	88.39	.8839E-01	98.28	1.72
3.75	74.33	.7433E-01	99.07	.93
4.00	62.50	.6250E-01	99.86	.14

## STATISTICS AND ENGINEERING PARAMETERS

SAMPLE D-1

## TINMAN'S STATISTICS:

MEDIAN 2.268 PHI 207.688 MICRONS  
 MEAN 2.092 PHI 234.492 MICRONS  
 STANDARD DEVIATION .708 MODERATELY WELL SORTED  
 SKEWNESS -.247 COARSE-SKewed  
 KURTOSIS 454 VERY PLATYKURTIC

FOLK'S STATISTICS:

MEAN 2.151 PHI 225.194 MICRONS  
 STANDARD DEVIATION .666 MODERATELY WELL SORTED  
 SKEWNESS -.281 COARSE-SKewed  
 KURTOSIS .890 PLATYKURTIC

## MOMENT MEASURES:

MEAN	2.140	PHI	226.887	MICRONS
SECOND MOMENT	.418			
STANDARD DEVIATION	.647	MODERATELY WELL SORTED		
THIRD MOMENT	-.262			
SKEWNESS	-.970	STRONGLY COARSE-SKewed		
FOURTH MOMENT	.733			
KURTOSIS	1.399	LEPTOKURTIC		

ENGINEERING PARAMETERS:

D60	2.111 PHI	231.538 MICRONS
D30	2.581 PHI	167.105 MICRONS
D10	2.895 PHI	134.452 MICRONS
COEF. OF UNIFORMITY	1.722	
COEF. OF CURVATURE	.897	

=====

TIERRA CONSULTING

SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

=====

ANALYSIS OF SAMPLE: D-2

PAN=.07=.1%

BVA-DIAL CORDY 2-26-07

SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 44.00

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.16	.36	99.64	.36
500.00	1.00	.89	2.02	97.61	2.39
250.00	2.00	7.08	16.09	81.52	18.48
125.00	3.00	33.69	76.57	4.95	95.05
62.50	4.00	2.11	4.80	.16	99.84

=====

TEXTURAL CLASSIFICATION AND PHI PERCENTILES                   SAMPLE D-2

=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF	2000.00 MICRONS
SAND/SILT CUTOFF	62.50 MICRONS
SILT/CLAY CUTOFF	4.00 MICRONS

PERCENT GRAVEL	.00
PERCENT SAND	99.84
PERCENT SILT AND CLAY	.16

## TEXTURAL DESCRIPTIONS

FOLK'S: SAND

SHEPARD'S: SAND

TREFETHEN'S: SAND

U.S. ARMY CORPS OF ENGINEERS: SAND

U. S. BUREAU OF STANDARDS

## PHI PERCENTILES

PHI(05)	1.16	PHI	446.76	MICRONS
PHI(16)	1.85	PHI	278.15	MICRONS
PHI(25)	2.09	PHI	235.67	MICRONS
PHI(50)	2.41	PHI	187.93	MICRONS
PHI(75)	2.74	PHI	149.87	MICRONS
PHI(84)	2.86	PHI	138.15	MICRONS
PHI(95)	3.00	PHI	125.05	MICRONS

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\* \*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

INTERPOLATED PHT PERCENTILES

SAMPLE D=2

PHI	MICRONS	MM	PCT.	RET	PCT. FINER
.00	1000.00	.1000E+01	.36		99.64
.25	840.90	.8409E+00	.87		99.13
.50	707.11	.7071E+00	1.38		98.63
.75	594.60	.5946E+00	1.88		98.12
1.00	500.00	.5000E+00	2.39		97.61
1.25	420.45	.4204E+00	6.41		93.59
1.50	353.55	.3536E+00	10.43		89.57
1.75	297.30	.2973E+00	14.45		85.55
2.00	250.00	.2500E+00	18.48		81.52
2.25	210.22	.2102E+00	37.62		62.38
2.50	176.78	.1768E+00	56.76		43.24
2.75	148.65	.1487E+00	75.90		24.10
3.00	125.00	.1250E+00	95.05		4.95
3.25	105.11	.1051E+00	96.24		3.76
3.50	88.39	.8839E-01	97.44		2.56
3.75	74.33	.7433E-01	98.64		1.36
4.00	62.50	.6250E-01	99.84		.16

=====

STATISTICS AND ENGINEERING PARAMETERS

SAMPLE D-2

=====

INMAN'S STATISTICS:

MEDIAN	2.412 PHI	187.935 MICRONS
MEAN	2.351 PHI	196.024 MICRONS
STANDARD DEVIATION	.505	MODERATELY WELL SORTED
SKEWNESS	-.120	COARSE-SKewed
KURTOSIS	.819	PLATyKURTIC

=====

FOLK'S STATISTICS:

MEAN	2.371 PHI	193.290 MICRONS
STANDARD DEVIATION	.531	MODERATELY WELL SORTED
SKEWNESS	-.240	COARSE-SKewed
KURTOSIS	1.153	LEPTOKURTIC

=====

MOMENT MEASURES:

MEAN	2.332 PHI	198.649 MICRONS
SECOND MOMENT	.295	
STANDARD DEVIATION	.544	MODERATELY WELL SORTED
THIRD MOMENT	-.219	
SKEWNESS	-1.366	STRONGLY COARSE-SKewed
FOURTH MOMENT	.628	
KURTOSIS	2.400	VERY LEPTOKURTIC

=====

ENGINEERING PARAMETERS:

D60	2.281 PHI	205.742 MICRONS
D30	2.673 PHI	156.811 MICRONS
D10	2.934 PHI	130.842 MICRONS
COEF. OF UNIFORMITY	1.572	
COEF. OF CURVATURE	.913	

=====

TIERRA CONSULTING  
SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

=====

ANALYSIS OF SAMPLE: D-3

PAN=0.02=.04%

BVA-DIAL CORDY 2-26-07

## SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 47.34

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.82	1.73	98.27	1.73
500.00	1.00	1.87	3.95	94.32	5.68
250.00	2.00	8.89	18.78	75.54	24.46
125.00	3.00	33.77	71.34	4.20	95.80
62.50	4.00	1.96	4.14	.06	99.94

===== TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE D-3 =====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
 SAND/SILT CUTOFF 62.50 MICRONS  
 SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
 PERCENT SAND 99.94

PERCENT SILT AND CLAY .06

## ===== TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
 SHEPARD'S: SAND  
 TREFETHEN'S: SAND  
 U.S. ARMY CORPS OF ENGINEERS: SAND  
 U.S. BUREAU OF SOILS: SAND

## ===== PHI PERCENTILES

PHI(05)	.83 PHI	563.59 MICRONS
PHI(16)	1.55 PHI	341.65 MICRONS
PHI(25)	2.01 PHI	248.69 MICRONS
PHI(50)	2.36 PHI	195.06 MICRONS
PHI(75)	2.71 PHI	152.99 MICRONS
PHI(84)	2.83 PHI	140.18 MICRONS
PHI(95)	2.99 PHI	125.97 MICRONS

===== EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

PHI	MICRONS	MM	PCT.	RET	PCT. FINER
.00	1000.00	.1000E+01	1.73		98.27
.25	840.90	.8409E+00	2.72		97.28
.50	707.11	.7071E+00	3.71		96.29
.75	594.60	.5946E+00	4.69		95.31
1.00	500.00	.5000E+00	5.68		94.32
1.25	420.45	.4204E+00	10.38		89.62
1.50	353.55	.3536E+00	15.07		84.93
1.75	297.30	.2973E+00	19.77		80.23
2.00	250.00	.2500E+00	24.46		75.54
2.25	210.22	.2102E+00	42.30		57.70
2.50	176.78	.1768E+00	60.13		39.87
2.75	148.65	.1487E+00	77.96		22.04
3.00	125.00	.1250E+00	95.80		4.20
3.25	105.11	.1051E+00	96.83		3.17
3.50	88.39	.8839E-01	97.87		2.13
3.75	74.33	.7433E-01	98.90		1.10
4.00	62.50	.6250E-01	99.94		.06

## STATISTICS AND ENGINEERING PARAMETERS

SAMPLE D-3

## TINMAN'S STATISTICS:

MEDIAN	2.358	PHI	195.060	MICRONS
MEAN	2.192	PHI	218.843	MICRONS
STANDARD DEVIATION	.643	MODERATELY WELL SORTED		
SKEWNESS	-.258	COARSE-SKewed		
KURTOSIS	.682	PLATYKURTIC		

FOLK'S STATISTICS:

MEAN 2.247 PHI 210.610 MICRONS  
 STANDARD DEVIATION .649 MODERATELY WELL SORTED  
 SKEWNESS -.337 STRONGLY COARSE-SKewed  
 KURTOSIS 1.264 LEPTOKURTIC

## MOMENT MEASURES:

MEAN	2.221	PHI	214.484	MICRONS
SECOND MOMENT	.466			
STANDARD DEVIATION	.683			MODERATELY WELL SORTED
THIRD MOMENT	-.519			
SKEWNESS	-1.630			STRONGLY COARSE-SKewed
FOURTH MOMENT	1.462			
KURTOSIS	2.243			VERY LEPTOKURTIC

ENGINEERING PARAMETERS:

D60	2.218 PHI	214.965 MICRONS
D30	2.638 PHI	160.609 MICRONS
D10	2.919 PHI	132.242 MICRONS
COEF. OF UNIFORMITY	1.626	
COEF. OF CURVATURE	.907	

=====  
TIERRA CONSULTING  
SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM  
=====

ANALYSIS OF SAMPLE: D-4

PAN=.01=.02%

BVA-DIAL CORDY 2-26-07

SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 44.00

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.13	.30	99.70	.30
500.00	1.00	1.84	4.18	95.52	4.48
250.00	2.00	22.62	51.41	44.11	55.89
125.00	3.00	18.28	41.55	2.57	97.43
62.50	4.00	.96	2.18	.39	99.61

=====  
TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE D-4  
=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF	2000.00 MICRONS
SAND/SILT CUTOFF	62.50 MICRONS
SILT/CLAY CUTOFF	4.00 MICRONS

PERCENT GRAVEL	.00
PERCENT SAND	99.61

PERCENT SILT AND CLAY	.39
-----------------------	-----

## TEXTURAL DESCRIPTIONS

FOLK'S: SAND

SHEPARD'S: SAND

TREFETHEN'S: SAND

U.S. ARMY CORPS OF ENGINEERS: SAND

U. S. BUREAU OF SOILS: SAND

## PHI PERCENTILES

PHI(05)	1.01	PHI	496.49	MICRONS
PHI(16)	1.22	PHI	428.05	MICRONS
PHI(25)	1.40	PHI	379.14	MICRONS
PHI(50)	1.89	PHI	270.65	MICRONS
PHI(75)	2.46	PHI	181.74	MICRONS
PHI(84)	2.68	PHI	156.40	MICRONS
PHI(95)	2.94	PHI	130.18	MICRONS

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\* \*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

## INTERPOLATED PHI PERCENTILES

SAMPLE D-4

PHI	MICRONS	MM	PCT.	RET	PCT. FINER
.00	1000.00	.1000E+01	.30		99.70
.25	840.90	.8409E+00	1.34		98.66
.50	707.11	.7071E+00	2.39		97.61
.75	594.60	.5946E+00	3.43		96.57
1.00	500.00	.5000E+00	4.48		95.52
1.25	420.45	.4204E+00	17.33		82.67
1.50	353.55	.3536E+00	30.18		69.82
1.75	297.30	.2973E+00	43.03		56.97
2.00	250.00	.2500E+00	55.89		44.11
2.25	210.22	.2102E+00	66.27		33.73
2.50	176.78	.1768E+00	76.66		23.34
2.75	148.65	.1487E+00	87.05		12.95
3.00	125.00	.1250E+00	97.43		2.57
3.25	105.11	.1051E+00	97.98		2.02
3.50	88.39	.8839E-01	98.52		1.48
3.75	74.33	.7433E-01	99.07		.93
4.00	62.50	.6250E-01	99.61		.39

## STATISTICS AND ENGINEERING PARAMETERS

SAMPLE D-4

## TINMAN'S STATISTICS:

MEDIAN 1.885 PHI 270.650 MICRONS  
 MEAN 1.950 PHI 258.741 MICRONS

STANDARD DEVIATION .726 MODERATELY SORTED  
 SKEWNESS .089 NEAR-SYMMETRICAL  
 KURTOSIS .330 VERY PLATYKURTIC

---

FOLK'S STATISTICS:

MEAN	1.929 PHI	262.652 MICRONS
STANDARD DEVIATION	.656	MODERATELY WELL SORTED
SKEWNESS	.091	NEAR-SYMMETRICAL
KURTOSIS	.746	PLATYKURTIC

---

MOMENT MEASURES:

MEAN	1.906 PHI	266.911 MICRONS
SECOND MOMENT	.387	
STANDARD DEVIATION	.622	MODERATELY WELL SORTED
THIRD MOMENT	-.016	
SKEWNESS	-.066	NEAR-SYMMETRICAL
FOURTH MOMENT	.469	
KURTOSIS	1.046	MESOKURTIC

---

ENGINEERING PARAMETERS:

D60	1.691 PHI	309.716 MICRONS
D30	2.340 PHI	197.549 MICRONS
D10	2.821 PHI	141.501 MICRONS
COEF. OF UNIFORMITY	2.189	
COEF. OF CURVATURE	.890	

---

TIERRA CONSULTING

SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

---

ANALYSIS OF SAMPLE: E-1

PAN=1.15=2.34%  
 BVA- DIAL CORDY 2-26-07

SIEVE ANALYSIS

---

SAMPLE WEIGHT = 48.95

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.26	.53	99.47	.53
500.00	1.00	.89	1.82	97.65	2.35
250.00	2.00	11.58	23.66	73.99	26.01
125.00	3.00	34.19	69.85	4.15	95.85
62.50	4.00	.84	1.72	2.43	97.57

=====  
TEXTURAL CLASSIFICATION AND PHI PERCENTILES                    SAMPLE E-1  
=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF        2000.00 MICRONS  
SAND/SILT CUTOFF          62.50 MICRONS  
SILT/CLAY CUTOFF          4.00 MICRONS

PERCENT GRAVEL            .00  
PERCENT SAND              97.57

PERCENT SILT AND CLAY    2.43

=====

TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
SHEPARD'S: SAND  
TREFETHEN'S: SAND  
U.S. ARMY CORPS OF ENGINEERS: SAND  
U.S. BUREAU OF SOILS: SAND

=====

PHI PERCENTILES

PHI(05)        1.11 PHI        462.64 MICRONS  
PHI(16)        1.58 PHI        335.17 MICRONS  
PHI(25)        1.96 PHI        257.48 MICRONS  
PHI(50)        2.34 PHI        197.03 MICRONS  
PHI(75)        2.70 PHI        153.74 MICRONS  
PHI(84)        2.83 PHI        140.60 MICRONS  
PHI(95)        2.99 PHI        126.06 MICRONS

=====

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*INSUFFICIENT DATA TO PERFORM EXTRAPOLATION\*\*\*

=====  
INTERPOLATED PHI PERCENTILES                    SAMPLE E-1  
=====

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	.53	99.47
.25	840.90	.8409E+00	.99	99.01
.50	707.11	.7071E+00	1.44	98.56
.75	594.60	.5946E+00	1.89	98.11
1.00	500.00	.5000E+00	2.35	97.65
1.25	420.45	.4204E+00	8.26	91.74
1.50	353.55	.3536E+00	14.18	85.82
1.75	297.30	.2973E+00	20.09	79.91
2.00	250.00	.2500E+00	26.01	73.99
2.25	210.22	.2102E+00	43.47	56.53

2.50	176.78	.1768E+00	60.93	39.07
2.75	148.65	.1487E+00	78.39	21.61
3.00	125.00	.1250E+00	95.85	4.15
3.25	105.11	.1051E+00	96.28	3.72
3.50	88.39	.8839E-01	96.71	3.29
3.75	74.33	.7433E-01	97.14	2.86
4.00	62.50	.6250E-01	97.57	2.43

-----  
STATISTICS AND ENGINEERING PARAMETERS                    SAMPLE E-1  
-----

## INMAN'S STATISTICS:

MEDIAN 2.344 PHI 197.029 MICRONS  
 MEAN 2.204 PHI 217.085 MICRONS  
 STANDARD DEVIATION .627 MODERATELY WELL SORTED  
 SKEWNESS -.223 COARSE-SKewed  
 KURTOSIS .497 VERY PLATykurtic

FOLK'S STATISTICS:

MEAN 2.250 PHI 210.183 MICRONS  
STANDARD DEVIATION .598 MODERATELY WELL SORTED  
SKEWNESS -.268 COARSE-SKewed  
KURTOSIS 1.033 MESOKURTIC

## MOMENT MEASURES:

\*\*INSUFFICIENT DATA FOR MOMENT MEASURES\*\*

## **ENGINEERING PARAMETERS:**

D60	2.200	PHI	217.585	MICRONS
D30	2.630	PHI	161.559	MICRONS
D10	2.916	PHI	132.475	MICRONS
COEF. OF UNIFORMITY	1.642			
COEF. OF CURVATURE	.906			

TIERRA CONSULTING

### SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

### ANALYSIS OF SAMPLE: E-2

PAN=.05=.1%  
BVA- DIAL CORDY 2-26-07

## SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 49.98

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.16	.32	99.68	.32
500.00	1.00	2.15	4.30	95.38	4.62
250.00	2.00	13.90	27.81	67.57	32.43
125.00	3.00	32.44	64.91	2.66	97.34
62.50	4.00	1.24	2.48	.18	99.82

===== TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE E-2 =====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
 SAND/SILT CUTOFF 62.50 MICRONS  
 SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
 PERCENT SAND 99.82  
 PERCENT SILT AND CLAY .18

## ===== TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
 SHEPARD'S: SAND  
 TREFETHEN'S: SAND  
 U.S. ARMY CORPS OF ENGINEERS: SAND  
 U.S. BUREAU OF SOILS: SAND

## ===== PHI PERCENTILES

PHI(05)	1.01 PHI	495.31 MICRONS
PHI(16)	1.41 PHI	376.54 MICRONS
PHI(25)	1.73 PHI	300.88 MICRONS
PHI(50)	2.27 PHI	207.24 MICRONS
PHI(75)	2.66 PHI	158.68 MICRONS
PHI(84)	2.79 PHI	144.14 MICRONS
PHI(95)	2.96 PHI	128.16 MICRONS

===== EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

PHI	MICRONS	MM	PCT.	RET	PCT. FINER
.00	1000.00	.1000E+01	.32		99.68
.25	840.90	.8409E+00	1.40		98.60
.50	707.11	.7071E+00	2.47		97.53
.75	594.60	.5946E+00	3.55		96.45
1.00	500.00	.5000E+00	4.62		95.38
1.25	420.45	.4204E+00	11.57		88.43
1.50	353.55	.3536E+00	18.53		81.47
1.75	297.30	.2973E+00	25.48		74.52
2.00	250.00	.2500E+00	32.43		67.57
2.25	210.22	.2102E+00	48.66		51.34
2.50	176.78	.1768E+00	64.89		35.11
2.75	148.65	.1487E+00	81.11		18.89
3.00	125.00	.1250E+00	97.34		2.66
3.25	105.11	.1051E+00	97.96		2.04
3.50	88.39	.8839E-01	98.58		1.42
3.75	74.33	.7433E-01	99.20		.80
4.00	62.50	.6250E-01	99.82		.18

=====  
STATISTICS AND ENGINEERING PARAMETERS                    SAMPLE E-2  
=====

## INMAN'S STATISTICS:

MEDIAN 2.271 PHI 207.236 MICRONS  
 MEAN 2.102 PHI 232.967 MICRONS  
 STANDARD DEVIATION .693 MODERATELY WELL SORTED  
 SKEWNESS -.244 COARSE-SKewed  
 KURTOSIS .408 VERY PLATykurtic

FOLK'S STATISTICS:

MEAN 2.158 PHI 224.053 MICRONS  
 STANDARD DEVIATION .642 MODERATELY WELL SORTED  
 SKEWNESS -.266 COARSE-SKewed  
 KURTOSIS .866 PLATykurtic

## MOMENT MEASURES:

MEAN	2.147	PHI	225.851	MICRONS
SECOND MOMENT	.382			
STANDARD DEVIATION	.618			MODERATELY WELL SORTED
THIRD MOMENT	-.236			
SKEWNESS	-1.002			STRONGLY COARSE-SKewed
FOURTH MOMENT	.615			
KURTOSIS	1.407			LEPTOKURTIC

ENGINEERING PARAMETERS:

D60	2.117 PHI	230.592 MICRONS
D30	2.579 PHI	167.381 MICRONS
D10	2.887 PHI	135.191 MICRONS
COEF. OF UNIFORMITY	1.706	
COEF. OF CURVATURE	.899	

=====

TIERRA CONSULTING  
SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

=====

ANALYSIS OF SAMPLE: E-3

PAN=.02=.04%  
BVA-DIAL CORDY 2-26-07

SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 43.54

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.01	.02	99.98	.02
500.00	1.00	2.86	6.57	93.41	6.59
250.00	2.00	27.82	63.90	29.51	70.49
125.00	3.00	12.44	28.57	.94	99.06
62.50	4.00	.39	.90	.05	99.95

=====

TEXTURAL CLASSIFICATION AND PHI PERCENTILES                    SAMPLE E-3

=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF	2000.00 MICRONS
SAND/SILT CUTOFF	62.50 MICRONS
SILT/CLAY CUTOFF	4.00 MICRONS
PERCENT GRAVEL	.00
PERCENT SAND	99.95

PERCENT SILT AND CLAY .05

=====

TEXTURAL DESCRIPTIONS

FOLK'S: SAND

SHEPARD'S: SAND

TREFETHEN'S: SAND

U.S. ARMY CORPS OF ENGINEERS: SAND

U.S. BUREAU OF SOILS: SAND

=====

PHI PERCENTILES

PHI(05)	.76 PHI	591.44 MICRONS
PHI(16)	1.15 PHI	451.49 MICRONS
PHI(25)	1.29 PHI	409.49 MICRONS
PHI(50)	1.68 PHI	312.22 MICRONS
PHI(75)	2.16 PHI	224.07 MICRONS
PHI(84)	2.47 PHI	180.12 MICRONS
PHI(95)	2.86 PHI	137.93 MICRONS

=====

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

=====

INTERPOLATED PHI PERCENTILES SAMPLE E-3

=====

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	.02	99.98
.25	840.90	.8409E+00	1.67	98.33
.50	707.11	.7071E+00	3.31	96.69
.75	594.60	.5946E+00	4.95	95.05
1.00	500.00	.5000E+00	6.59	93.41
1.25	420.45	.4204E+00	22.57	77.43
1.50	353.55	.3536E+00	38.54	61.46
1.75	297.30	.2973E+00	54.51	45.49
2.00	250.00	.2500E+00	70.49	29.51
2.25	210.22	.2102E+00	77.63	22.37
2.50	176.78	.1768E+00	84.77	15.23
2.75	148.65	.1487E+00	91.92	8.08
3.00	125.00	.1250E+00	99.06	.94
3.25	105.11	.1051E+00	99.28	.72
3.50	88.39	.8839E-01	99.51	.49
3.75	74.33	.7433E-01	99.73	.27
4.00	62.50	.6250E-01	99.95	.05

=====

STATISTICS AND ENGINEERING PARAMETERS SAMPLE E-3

=====

INMAN'S STATISTICS:

MEDIAN	1.679 PHI	312.220 MICRONS
MEAN	1.810 PHI	285.171 MICRONS
STANDARD DEVIATION	.663	MODERATELY WELL SORTED
SKEWNESS	.197	FINE-SKewed
KURTOSIS	.584	VERY PLATYKURTIC

=====

FOLK'S STATISTICS:

MEAN	1.767 PHI	293.916 MICRONS
STANDARD DEVIATION	.650	MODERATELY WELL SORTED
SKEWNESS	.160	FINE-SKewed
KURTOSIS	.990	MESOKURTIC

---

MOMENT MEASURES:

MEAN	1.737 PHI	300.036 MICRONS
SECOND MOMENT	.332	
STANDARD DEVIATION	.576	MODERATELY WELL SORTED
THIRD MOMENT	.041	
SKEWNESS	.214	FINE-SKewed
FOURTH MOMENT	.345	
KURTOSIS	1.045	MESOKURTIC

---

ENGINEERING PARAMETERS:

D60	1.523 PHI	347.995 MICRONS
D30	1.992 PHI	251.324 MICRONS
D10	2.683 PHI	155.722 MICRONS
COEF. OF UNIFORMITY	2.235	
COEF. OF CURVATURE	1.166	

---

TIERRA CONSULTING

SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

---

ANALYSIS OF SAMPLE: E-4

PAN=.02=.04%

BVA DIAL CORDY 2-26-07

SIEVE ANALYSIS

---

SAMPLE WEIGHT = 48.72

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.06	.12	99.88	.12
500.00	1.00	2.71	5.56	94.31	5.69
250.00	2.00	27.05	55.52	38.79	61.21
125.00	3.00	18.26	37.48	1.31	98.69
62.50	4.00	.55	1.13	.18	99.82

TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE E-4

SAMPLE E-4

**SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION**

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
 SAND/SILT CUTOFF 62.50 MICRONS  
 SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
PERCENT SAND 99.82  
PERCENT SILT AND CLAY .18

## TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
SHEPARD'S: SAND  
TREFETHEN'S: SAND  
U.S. ARMY CORPS OF ENGINEERS: SAND  
U.S. BUREAU OF SOILS: SAND

## PHT PERCENTILES

PHI(05)	.88	PHI	544.59	MICRONS
PHI(16)	1.19	PHI	439.59	MICRONS
PHI(25)	1.35	PHI	392.87	MICRONS
PHI(50)	1.80	PHI	287.54	MICRONS
PHI(75)	2.37	PHI	193.71	MICRONS
PHI(84)	2.61	PHI	164.01	MICRONS
PHI(95)	2.90	PHI	133.82	MICRONS

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

INTERPOLATED PHT PERCENTILES SAMPLE E-4

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	.12	99.88
.25	840.90	.8409E+00	1.51	98.49
.50	707.11	.7071E+00	2.90	97.10
.75	594.60	.5946E+00	4.29	95.71
1.00	500.00	.5000E+00	5.69	94.31
1.25	420.45	.4204E+00	19.57	80.43
1.50	353.55	.3536E+00	33.45	66.55
1.75	297.30	.2973E+00	47.33	52.67
2.00	250.00	.2500E+00	61.21	38.79
2.25	210.22	.2102E+00	70.58	29.42
2.50	176.78	.1768E+00	79.95	20.05
2.75	148.65	.1487E+00	89.32	10.68
3.00	125.00	.1250E+00	98.69	1.31

3.25	105.11	.1051E+00	98.97	1.03
3.50	88.39	.8839E-01	99.25	.75
3.75	74.33	.7433E-01	99.53	.47
4.00	62.50	.6250E-01	99.82	.18

## INMAN'S STATISTICS:

MEDIAN 1.798 PHI 287.543 MICRONS  
 MEAN 1.897 PHI 268.508 MICRONS  
 STANDARD DEVIATION .711 MODERATELY SORTED  
 SKEWNESS .139 FINE-SKewed  
 KURTOSIS .424 VERY PLATykURTIC

FOLK'S STATISTICS:

MEAN 1.864 PHI 274.709 MICRONS  
STANDARD DEVIATION .662 MODERATELY WELL SORTED  
SKEWNESS .114 FINE-SKewed  
KURTOSIS .813 PLATykurtic

## MOMENT MEASURES:

MEAN	1.837	PHI	279.997	MICRONS
SECOND MOMENT	.365			
STANDARD DEVIATION	.604	MODERATELY WELL SORTED		
THIRD MOMENT	-.008			
SKEWNESS	-.037	NEAR-SYMMETRICAL		
FOURTH MOMENT	.380			
KURTOSIS	.951	MESOKURTIC		

#### ENGINEERING PARAMETERS:

D60	1.618	PHI	325.778	MICRONS
D30	2.235	PHI	212.479	MICRONS
D10	2.768	PHI	146.784	MICRONS
COEF. OF UNIFORMITY	2.219			
COEF. OF CURVATURE	.944			

TIERRA CONSULTING

SEDIMENTARY PETROLOGY ANALYSTS PROGRAM

ANALYSIS OF SAMPLE: F-1

$$PAN = 0.10 = .2\%$$

BVA DIAL CORDY 2-26-07

## SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 44.93

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.13	.29	99.71	.29
500.00	1.00	2.16	4.81	94.90	5.10
250.00	2.00	14.22	31.65	63.25	36.75
125.00	3.00	27.48	61.16	2.09	97.91
62.50	4.00	.79	1.76	.33	99.67

## ===== TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE F-1 =====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF	2000.00 MICRONS
SAND/SILT CUTOFF	62.50 MICRONS
SILT/CLAY CUTOFF	4.00 MICRONS
PERCENT GRAVEL	.00
PERCENT SAND	99.67

PERCENT SILT AND CLAY .33

## ===== TEXTURAL DESCRIPTIONS =====

FOLK'S: SAND  
 SHEPARD'S: SAND  
 TREFETHEN'S: SAND  
 U.S. ARMY CORPS OF ENGINEERS: SAND  
 U.S. BUREAU OF SOILS: SAND

## ===== PHI PERCENTILES =====

PHI(05)	.98 PHI	507.03 MICRONS
PHI(16)	1.34 PHI	393.79 MICRONS
PHI(25)	1.63 PHI	323.34 MICRONS
PHI(50)	2.22 PHI	215.13 MICRONS
PHI(75)	2.63 PHI	162.05 MICRONS
PHI(84)	2.77 PHI	146.34 MICRONS
PHI(95)	2.95 PHI	129.19 MICRONS

===== EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	.29	99.71
.25	840.90	.8409E+00	1.49	98.51
.50	707.11	.7071E+00	2.69	97.31
.75	594.60	.5946E+00	3.89	96.11
1.00	500.00	.5000E+00	5.10	94.90
1.25	420.45	.4204E+00	13.01	86.99
1.50	353.55	.3536E+00	20.92	79.08
1.75	297.30	.2973E+00	28.83	71.17
2.00	250.00	.2500E+00	36.75	63.25
2.25	210.22	.2102E+00	52.04	47.96
2.50	176.78	.1768E+00	67.33	32.67
2.75	148.65	.1487E+00	82.62	17.38
3.00	125.00	.1250E+00	97.91	2.09
3.25	105.11	.1051E+00	98.35	1.65
3.50	88.39	.8839E-01	98.79	1.21
3.75	74.33	.7433E-01	99.23	.77
4.00	62.50	.6250E-01	99.67	.33

## INMAN'S STATISTICS:

MEDIAN	2.217	PHI	215.132	MICRONS
MEAN	2.059	PHI	240.057	MICRONS
STANDARD DEVIATION	.714	MODERATELY SORTED		
SKEWNESS	-.221	COARSE-SKewed		
KURTOSIS	.381	VERY PLATykurtic		

FOLK'S STATISTICS:

MEAN	2.111	PHI	231.443	MICRONS
STANDARD DEVIATION	.656	MODERATELY WELL SORTED		
SKEWNESS	-.238	COARSE-SKewed		
KURTOSIS	.811	PLATYKURTIC		

---

## MOMENT MEASURES:

MEAN	2.088	PHI	235.220	MICRONS
SECOND MOMENT	.389			
STANDARD DEVIATION	.624	MODERATELY WELL SORTED		
THIRD MOMENT	-.215			
SKEWNESS	-.885	STRONGLY COARSE-SKewed		
FOURTH MOMENT	.561			
KURTOSIS	1.236	LEPTOKURTIC		

=====

ENGINEERING PARAMETERS:

D60	2.053 PHI	240.949 MICRONS
D30	2.544 PHI	171.502 MICRONS
D10	2.871 PHI	136.720 MICRONS
COEF. OF UNIFORMITY	1.762	
COEF. OF CURVATURE	.893	

=====

TIERRA CONSULTING  
SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

=====

ANALYSIS OF SAMPLE: F-2

PAN=.05=.11%  
BVA DIAL CORDY 2-26-07

SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 44.73

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	9.78	21.86	78.14	21.86
500.00	1.00	19.82	44.31	33.83	66.17
250.00	2.00	9.72	21.73	12.09	87.91
125.00	3.00	5.13	11.47	.63	99.37
62.50	4.00	.18	.40	.22	99.78

=====

TEXTURAL CLASSIFICATION AND PHI PERCENTILES                    SAMPLE F-2

=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF	2000.00 MICRONS
SAND/SILT CUTOFF	62.50 MICRONS
SILT/CLAY CUTOFF	4.00 MICRONS

PERCENT GRAVEL	.00
PERCENT SAND	99.78
PERCENT SILT AND CLAY	.22

## TEXTURAL DESCRIPTIONS

FOLK'S: SAND

SHEPARD'S: SAND

TREFETHEN'S: SAND

U.S. ARMY CORPS OF ENGINEERS: SAND

U. S. BUREAU OF STANDARDS

## PHI PERCENTILES

PHI( 25)	.07	PHI	952.14	MICRONS
PHI( 50)	.63	PHI	643.96	MICRONS
PHI( 75)	1.41	PHI	377.32	MICRONS
PHI( 84)	1.82	PHI	283.16	MICRONS
PHI( 95)	2.62	PHI	162.82	MICRONS

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

## INTERPOLATED PHI PERCENTILES

SAMPLE F-2

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	21.86	78.14
.25	840.90	.8409E+00	32.94	67.06
.50	707.11	.7071E+00	44.02	55.98
.75	594.60	.5946E+00	55.10	44.90
1.00	500.00	.5000E+00	66.17	33.83
1.25	420.45	.4204E+00	71.61	28.39
1.50	353.55	.3536E+00	77.04	22.96
1.75	297.30	.2973E+00	82.47	17.53
2.00	250.00	.2500E+00	87.91	12.09
2.25	210.22	.2102E+00	90.77	9.23
2.50	176.78	.1768E+00	93.64	6.36
2.75	148.65	.1487E+00	96.51	3.49
3.00	125.00	.1250E+00	99.37	.63
3.25	105.11	.1051E+00	99.47	.53
3.50	88.39	.8839E-01	99.58	.42
3.75	74.33	.7433E-01	99.68	.32
4.00	62.50	.6250E-01	99.78	.22

## STATISTICS AND ENGINEERING PARAMETERS

SAMPLE F-2

## TINMAN'S STATISTICS:

MEDIAN .635 PHT 643.957 MICRONS

FOLK'S STATISTICS:

\*\*INSUFFICIENT DATA FOR FOLK'S STATISTICS\*\*

---

MOMENT MEASURES:

\*\*WARNING\*\* ACCURACY OF MOMENT MEASURES MAY BE  
ADVERSELY AFFECTED BY THE LARGE AMOUNT OF  
MATERIAL RETAINED IN THE FIRST SIEVE \*\*

MEAN	.739 PHI	599.159 MICRONS
SECOND MOMENT	.873	
STANDARD DEVIATION	.934	MODERATELY SORTED
THIRD MOMENT	.385	
SKEWNESS	.472	STRONGLY FINE-SKEWED
FOURTH MOMENT	1.926	
KURTOSIS	.842	PLATYKURTIC

---

ENGINEERING PARAMETERS:

D60	.409 PHI	752.997 MICRONS
D30	1.176 PHI	442.568 MICRONS
D10	2.183 PHI	220.271 MICRONS
COEF. OF UNIFORMITY	3.419	
COEF. OF CURVATURE	1.181	

---

TIERRA CONSULTING

SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

---

ANALYSIS OF SAMPLE: F-3

PAN=0.04=.08%

SIEVE ANALYSIS

---

SAMPLE WEIGHT = 46.34

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.19	.41	99.59	.41
500.00	1.00	1.43	3.09	96.50	3.50
250.00	2.00	8.47	18.28	78.23	21.77
125.00	3.00	34.92	75.36	2.87	97.13
62.50	4.00	1.26	2.72	.15	99.85

=====  
TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE F-3  
=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
SAND/SILT CUTOFF 62.50 MICRONS  
SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
PERCENT SAND 99.85

PERCENT SILT AND CLAY .15

=====

TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
SHEPARD'S: SAND  
TREFETHEN'S: SAND  
U.S. ARMY CORPS OF ENGINEERS: SAND  
U.S. BUREAU OF SOILS: SAND

=====

PHI PERCENTILES

PHI(05)	1.08	PHI	472.28	MICRONS
PHI(16)	1.68	PHI	311.20	MICRONS
PHI(25)	2.04	PHI	242.69	MICRONS
PHI(50)	2.37	PHI	192.83	MICRONS
PHI(75)	2.71	PHI	153.22	MICRONS
PHI(84)	2.83	PHI	141.05	MICRONS
PHI(95)	2.97	PHI	127.47	MICRONS

=====

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

=====  
INTERPOLATED PHI PERCENTILES SAMPLE F-3  
=====

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	.41	99.59
.25	840.90	.8409E+00	1.18	98.82
.50	707.11	.7071E+00	1.95	98.05
.75	594.60	.5946E+00	2.72	97.28
1.00	500.00	.5000E+00	3.50	96.50
1.25	420.45	.4204E+00	8.07	91.93
1.50	353.55	.3536E+00	12.63	87.37

1.75	297.30	.2973E+00	17.20	82.80
2.00	250.00	.2500E+00	21.77	78.23
2.25	210.22	.2102E+00	40.61	59.39
2.50	176.78	.1768E+00	59.45	40.55
2.75	148.65	.1487E+00	78.29	21.71
3.00	125.00	.1250E+00	97.13	2.87
3.25	105.11	.1051E+00	97.81	2.19
3.50	88.39	.8839E-01	98.49	1.51
3.75	74.33	.7433E-01	99.17	.83
4.00	62.50	.6250E-01	99.85	.15

## STATISTICS AND ENGINEERING PARAMETERS

SAMPLE F-3

## INMAN'S STATISTICS:

MEDIAN	2.375	PHI	192.834	MICRONS
MEAN	2.255	PHI	209.506	MICRONS
STANDARD DEVIATION	.571	MODERATELY WELL SORTED		
SKEWNESS	-.210	COARSE-SKewed		
KURTOSIS	.655	VERY PLATykurtic		

## FOLK'S STATISTICS:

MEAN 2.295 PHI 203.794 MICRONS  
 STANDARD DEVIATION .572 MODERATELY WELL SORTED  
 SKEWNESS -.289 COARSE-SKewed  
 KURTOSIS 1.167 LEPTOKURTIC

## MOMENT MEASURES:

MEAN	2.267	PHI	207.817	MICRONS
SECOND MOMENT	.318			
STANDARD DEVIATION	.563			MODERATELY WELL SORTED
THIRD MOMENT	-.279			
SKEWNESS	-1.558			STRONGLY COARSE-SKewed
FOURTH MOMENT	.669			
KURTOSIS	2.212			VERY LEPTOKURTIC

## ENGINEERING PARAMETERS:

D60	2.242	PHI	211.413	MICRONS
D30	2.640	PHI	160.431	MICRONS
D10	2.905	PHI	133.473	MICRONS
COEF. OF UNIFORMITY	1.584			
COEF. OF CURVATURE	.912			

TIERRA CONSULTING

## SEDPET III: SEDIMENTARY PETROLOGY ANALYSIS PROGRAM

ANALYSIS OF SAMPLE: F-4

PAN=.03=.06%

BVA-DIAL CORDY 2-26-07

SIEVE ANALYSIS

=====

SAMPLE WEIGHT = 44.28

SIEVE SIZE (MICRONS)	SIEVE SIZE (PHI)	SIEVE WT (GRAMS)	WEIGHT PERCENT RETAINED	CUMULATIVE PERCENT THROUGH	CUMULATIVE PERCENT RETAINED
1000.00	.00	.13	.29	99.71	.29
500.00	1.00	1.41	3.18	96.52	3.48
250.00	2.00	9.70	21.91	74.62	25.38
125.00	3.00	32.14	72.58	2.03	97.97
62.50	4.00	.86	1.94	.09	99.91

=====

TEXTURAL CLASSIFICATION AND PHI PERCENTILES SAMPLE F-4

=====

SIZE CLASSIFICATION USED: STANDARD GEOLOGICAL SOILS CLASSIFICATION

GRAVEL/SAND CUTOFF 2000.00 MICRONS  
SAND/SILT CUTOFF 62.50 MICRONS  
SILT/CLAY CUTOFF 4.00 MICRONS

PERCENT GRAVEL .00  
PERCENT SAND 99.91

PERCENT SILT AND CLAY .09

=====

TEXTURAL DESCRIPTIONS

FOLK'S: SAND  
SHEPARD'S: SAND  
TREFETHEN'S: SAND  
U.S. ARMY CORPS OF ENGINEERS: SAND  
U.S. BUREAU OF SOILS: SAND

=====

PHI PERCENTILES

PHI(05)	1.07 PHI	476.49 MICRONS
PHI(16)	1.57 PHI	336.43 MICRONS
PHI(25)	1.98 PHI	253.06 MICRONS
PHI(50)	2.34 PHI	197.63 MICRONS
PHI(75)	2.68 PHI	155.66 MICRONS

PHI(84) 2.81 PHI 142.84 MICRONS  
PHI(95) 2.96 PHI 128.59 MICRONS

=====

EXTRAPOLATION OF DATA TO CUMULATIVE 99.5% RETAINED.

\*\*\*OVER 99.5% SAMPLE ACCOUNTED FOR BY INPUT DATA - NO EXTRAPOLATION REQUIRED.

=====  
INTERPOLATED PHI PERCENTILES SAMPLE F-4  
=====

PHI	MICRONS	MM	PCT. RET	PCT. FINER
.00	1000.00	.1000E+01	.29	99.71
.25	840.90	.8409E+00	1.09	98.91
.50	707.11	.7071E+00	1.89	98.11
.75	594.60	.5946E+00	2.68	97.32
1.00	500.00	.5000E+00	3.48	96.52
1.25	420.45	.4204E+00	8.95	91.05
1.50	353.55	.3536E+00	14.43	85.57
1.75	297.30	.2973E+00	19.91	80.09
2.00	250.00	.2500E+00	25.38	74.62
2.25	210.22	.2102E+00	43.53	56.47
2.50	176.78	.1768E+00	61.68	38.32
2.75	148.65	.1487E+00	79.82	20.18
3.00	125.00	.1250E+00	97.97	2.03
3.25	105.11	.1051E+00	98.45	1.55
3.50	88.39	.8839E-01	98.94	1.06
3.75	74.33	.7433E-01	99.42	.58
4.00	62.50	.6250E-01	99.91	.09

=====  
STATISTICS AND ENGINEERING PARAMETERS SAMPLE F-4  
=====

INMAN'S STATISTICS:

MEDIAN 2.339 PHI 197.628 MICRONS  
MEAN 2.190 PHI 219.213 MICRONS  
STANDARD DEVIATION .618 MODERATELY WELL SORTED  
SKEWNESS -.242 COARSE-SKewed  
KURTOSIS .529 VERY PLATYKURTIC

=====

FOLK'S STATISTICS:

MEAN 2.239 PHI 211.768 MICRONS  
STANDARD DEVIATION .595 MODERATELY WELL SORTED  
SKEWNESS -.293 COARSE-SKewed  
KURTOSIS 1.105 MESOKURTIC

=====

MOMENT MEASURES:

MEAN	2.226 PHI	213.808 MICRONS
SECOND MOMENT	.318	
STANDARD DEVIATION	.564	MODERATELY WELL SORTED
THIRD MOMENT	-.252	
SKEWNESS	-1.402	STRONGLY COARSE-SKewed
FOURTH MOMENT	.560	
KURTOSIS	1.846	VERY LEPTOKURTIC

---

ENGINEERING PARAMETERS:

D60	2.201 PHI	217.431 MICRONS
D30	2.615 PHI	163.268 MICRONS
D10	2.890 PHI	134.882 MICRONS
COEF. OF UNIFORMITY	1.612	
COEF. OF CURVATURE	.909	

---

**APPENDIX C**

**Station Data Summary Report**

## Station Data Summary Report

### Station A1

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: A1  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Oweniida					
Oweniidae					
<i>Owenia fusiformis</i>	1	25	1	14.3	25
Spionida					
Chaetopteridae					
<i>Spiochaetopterus oculatus</i>	1	25	1	14.3	25
Cirratulidae					
<i>Caulieriella sp. J</i>	1	25	1	14.3	25
Magelonidae					
<i>Magelona (LPIL)</i>	2	50	2	28.6	50
Terebellida					
Pectinariidae					
<i>Pectinaria gouldii</i>	1	25	1	14.3	25
<b>Mollusca</b>					
Bivalvia					
<i>Bivalvia (LPIL)</i>	1	25	1	14.3	25

Note LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

## Station Data Summary Report

### Station A1

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 2

Station: A1  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

### DATA SUMMARY

FAUNAL PARAMETERS	Station	Rep 1	Mean	Std Dev
Total Taxa	6	6	6	0
Total Individuals	7	7	7	0
Density (nos/sq.m.)		175	175	0

### FAUNAL INDICES

Species Diversity (Shannon; log base e)	H' = 1.75
Species Diversity (Shannon; log base 2)	d = 2.52
Species Diversity (Shannon; log base 10)	H = 0.76
Species Diversity (Simpson; 1/S)	1/S = 21
Species Evenness (Pielou)	J' = 0.98
Species Richness (Margalef)	D = 2.57
Equitability Index (Lloyd & Ghelardi)	e = 1.33

MAJOR TAXONOMIC GROUPS	Total No. Taxa	Taxa % Total	Total No. Individuals	Individuals % Total
Annelida	5	83.3	6	85.7
Mollusca	1	16.6	1	14.2
<b>TOTALS</b>	<b>6</b>		<b>7</b>	

### Station Data Summary Report

#### Station A2

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 2

Station: A2  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
<b>Polychaeta</b>					
<b>Capitellida</b>					
<b>Capitellidae</b>					
Capitellidae (LPIL)	1	25	1	5.9	25
<b>Opheliida</b>					
<b>Opheliidae</b>					
Armandia maculata	1	25	1	5.9	25
<b>Phyllodocida</b>					
<b>Glyceridae</b>					
Glyceridae (LPIL)	2	50	2	11.8	50
<b>Nephtyidae</b>					
Aglaophamus verrilli	1	25	1	5.9	25
<b>Spionida</b>					
<b>Cirratulidae</b>					
Caulleriella sp. J	2	50	2	11.8	50
<b>Magelonidae</b>					
Magelona (LPIL)	1	25	1	5.9	25
<b>Mollusca</b>					
<b>Bivalvia</b>					
<b>Veneroida</b>					
<b>Lucinidae</b>					
Lucinidae (LPIL)	2	50	2	11.8	50
<b>Tellinidae</b>					
Tellina (LPIL)	2	50	2	11.8	50
Tellina iris	1	25	1	5.9	25

**Station Data Summary Report****Station A2**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 2

Station: A2  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Gastropoda					
Cephalaspidea					
Scaphandridae					
Cylichna alba	4	100	4	23.5	100

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station A2**

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 3

Station: A2  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	10	10	10	0
Total Individuals	17	17	17	0
Density (nos/sq.m.)		425	425	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 2.18
Species Diversity (Shannon; log base 2)	d = 3.15
Species Diversity (Shannon; log base 10)	H = 0.95
Species Diversity (Simpson; 1/S)	1/S = 13.6
Species Evenness (Pielou)	J' = 0.95
Species Richness (Margalef)	D = 3.18
Equitability Index (Lloyd & Ghelardi)	e = 1.25

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Annelida	6	60	8	47
Mollusca	4	40	9	52.9
<b>TOTALS</b>	<b>10</b>		<b>17</b>	

### Station Data Summary Report

#### Station A3

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: A3  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
<b>Spionida</b>					
Magelonidae					
<i>Magelona papillicornis</i>	2	50	2	20	50
Spionidae					
<i>Parapriionospio pinnata</i>	1	25	1	10	25
<i>Spionidae (LPIL)</i>	1	25	1	10	25
<b>Arthropoda</b>					
Ostracoda					
Myodocopina					
Sarsiellidae					
<i>Eusarsiella (LPIL)</i>	1	25	1	10	25
<b>Cnidaria</b>					
Hydrozoa					
<i>Hydrozoa (LPIL)</i>	1	25	1	10	25
<b>Mollusca</b>					
Gastropoda					
Cephalaspidea					
Scaphandridae					
<i>Acteocina canaliculata</i>	1	25	1	10	25
<i>Cylichna alba</i>	1	25	1	10	25

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

### Station Data Summary Report

#### Station A3

Page 1

Station: A3

Sample Type: Macrofauna

Replicates: 1

Sample Area: 0.0400

#### TAXON

	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Phoronida					
Phoronidae					
Phoronis (LPIL)	1	25	1	10	25
Rhynchocoela					
Rhynchocoela (LPIL)	1	25	1	10	25

Note:

LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station A3**

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 3

Station: A3  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	9	9	9	0
Total Individuals	10	10	10	0
Density (nos/sq.m.)	250	250	0	

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' =	2.16
Species Diversity (Shannon; log base 2)	d =	3.12
Species Diversity (Shannon; log base 10)	H =	0.94
Species Diversity (Simpson; 1/S)	1/S =	45
Species Evenness (Pielou)	J' =	0.98
Species Richness (Margalef)	D =	3.47
Equitability Index (Lloyd & Ghelardi)	e =	1.37

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa % Total</b>	<b>Total No.</b>	<b>Individuals % Total</b>
	Taxa	% Total	Individuals	% Total
Cnidaria	1	11.1	1	10
Rhynchocoela	1	11.1	1	10
Annelida	3	33.3	4	40
Mollusca	2	22.2	2	20
Arthropoda	1	11.1	1	10
Phoronida	1	11.1	1	10
<b>TOTALS</b>	<b>9</b>		<b>10</b>	

### Station Data Summary Report

#### Station A4

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: A4  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Capitellida					
Capitellidae					
<i>Mediomastus californiensis</i>	1	25	1	9.1	25
Eunicida					
Onuphidae					
<i>Onuphidae (LPIL)</i>	1	25	1	9.1	25
Phyllodocida					
Goniadidae					
<i>Goniada littorea</i>	1	25	1	9.1	25
Spionida					
Cirratulidae					
<i>Cauilleriella sp. J</i>	1	25	1	9.1	25
<b>Arthropoda</b>					
Malacostraca					
Amphipoda					
Platyischnopidae					
<i>Eudevenopus honduranus</i>	1	25	1	9.1	25
Synopiidae					
Synopiidae (LPIL)	2	50	2	18.2	50
<b>Mollusca</b>					
Bivalvia					
<i>Bivalvia (LPIL)</i>	1	25	1	9.1	25

**Station Data Summary Report****Station A4**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: A4  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Veneroida					
Mesodesmatidae					
<i>Ervilia concentrica</i>	1	25	1	9.1	25
Gastropoda					
Neogastropoda					
Olividae					
<i>Oliva sayana</i>	2	50	2	18.2	50

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

## Station Data Summary Report

### Station A4

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 3

Station: A4  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

### DATA SUMMARY

FAUNAL PARAMETERS	Station	Rep 1	Mean	Std Dev
Total Taxa	9	9	9	0
Total Individuals	11	11	11	0
Density (nos/sq.m.)		275	275	0

### FAUNAL INDICES

Species Diversity (Shannon; log base e)	H' = 2.15
Species Diversity (Shannon; log base 2)	d = 3.1
Species Diversity (Shannon; log base 10)	H = 0.93
Species Diversity (Simpson; 1/S)	1/S = 27.5
Species Evenness (Pielou)	J' = 0.98
Species Richness (Margalef)	D = 3.34
Equitability Index (Lloyd & Ghelardi)	e = 1.34

MAJOR TAXONOMIC GROUPS	Total No.	Taxa	Total No.	Individuals
	Taxa	% Total	Individuals	% Total
Annelida	4	44.4	4	36.3
Mollusca	3	33.3	4	36.3
Arthropoda	2	22.2	3	27.2
<b>TOTALS</b>	<b>9</b>		<b>11</b>	

### Station Data Summary Report

#### Station A5

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: A5  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Orbiniida					
Paraonidae					
Aricidea catherinae	1	25	1	9.1	25
Phyllodocida					
Pilargiidae					
Sigambra tentaculata	1	25	1	9.1	25
Spionida					
Magelonidae					
Magelona papillicornis	3	75	3	27.3	75
<b>Arthropoda</b>					
Ostracoda					
Myodocopina					
Sarsiellidae					
Eusarsiella texana	1	25	1	9.1	25
<b>Mollusca</b>					
Bivalvia					
Veneroida					
Tellinidae					
Tellina (LPIL)	3	75	3	27.3	75
Gastropoda					
Neogastropoda					
Olividae					
Oliva sayana	1	25	1	9.1	25

**Station Data Summary Report****Station A5**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: A5  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Pyramidelloida					
Pyramidellidae					
<i>Odostomia weberi</i>	1	25	1	9.1	25

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

## Station Data Summary Report

### Station A5

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 3

Station: A5  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

#### **DATA SUMMARY**

FAUNAL PARAMETERS	Station	Rep 1	Mean	Std Dev
Total Taxa	7	7	7	0
Total Individuals	11	11	11	0
Density (nos/sq.m.)		275	275	0

#### **FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' =	1.8
Species Diversity (Shannon; log base 2)	d =	2.59
Species Diversity (Shannon; log base 10)	H =	0.78
Species Diversity (Simpson; 1/S)	1/S =	9.17
Species Evenness (Pielou)	J' =	0.92
Species Richness (Margalef)	D =	2.5
Equitability Index (Lloyd & Ghelardi)	e =	1.2

MAJOR TAXONOMIC GROUPS	Total No.	Taxa	Total No.	Individuals
	Taxa	% Total	Individuals	% Total
Annelida	3	42.8	5	45.4
Mollusca	3	42.8	5	45.4
Arthropoda	1	14.2	1	9
<b>TOTALS</b>	<b>7</b>		<b>11</b>	

## Station Data Summary Report

### Station B1

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: B7  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Capitellida					
Capitellidae					
<i>Mediomastus (LPIL)</i>	3	75	3	12	75
Eunicida					
Dorvilleidae					
<i>Schistomeringos pectinata</i>	1	25	1	4	25
Phyllodocida					
Chrysopetalidae					
<i>Bhawania heteroseta</i>	2	50	2	8	50
Goniadidae					
<i>Goniadides caroliniae</i>	2	50	2	8	50
Spionida					
Cirratulidae					
<i>Cirratulidae (LPIL)</i>	1	25	1	4	25
<b>Arthropoda</b>					
Malacostraca					
Amphipoda					
Aoridae					
<i>Rildardanus laminosa</i>	1	25	1	4	25
Melitidae					
<i>Maera caroliniana</i>	2	50	2	8	50
Synopiidae					
<i>Metatiron triocellatus</i>	1	25	1	4	25

### Station Data Summary Report

#### Station B1

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: B7  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Echinodermata					
Ophiuroidea					
Ophiuroidea (LPIL)	2	50	2	8	50
Ophiurida					
Amphiuridae					
Amphiuridae (LPIL)	1	25	1	4	25
Mollusca					
Bivalvia					
Veneroida					
Crassatellidae					
Crassinella lunulata	4	100	4	16	100
Gastropoda					
Mesogastropoda					
Caecidae					
Caecum pulchellum	1	25	1	4	25
Platyhelminthes					
Turbellaria					
Turbellaria (LPIL)	2	50	2	8	50
Rhynchocoela					
Rhynchocoela (LPIL)	2	50	2	8	50

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station B1**

Page 3

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Station: B7  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	14	14	14	0
Total Individuals	25	25	25	0
Density (nos/sq.m.)		625	625	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' =	2.53
Species Diversity (Shannon; log base 2)	d =	3.65
Species Diversity (Shannon; log base 10)	H =	1.1
Species Diversity (Simpson; 1/S)	1/S =	20
Species Evenness (Pielou)	J' =	0.96
Species Richness (Margalef)	D =	4.04
Equitability Index (Lloyd & Ghelardi)	e =	1.29

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Platyhelminthes	1	7.1	2	8
Rhynchocoela	1	7.1	2	8
Annelida	5	35.7	9	36
Mollusca	2	14.2	5	20
Arthropoda	3	21.4	4	16
Echinodermata	2	14.2	3	12
<b>TOTALS</b>	<b>14</b>		<b>25</b>	

### Station Data Summary Report

#### Station B2

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: B2  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
<b>Polychaeta</b>					
<b>Capitellida</b>					
<b>Capitellidae</b>					
<i>Mediomastus (LPIL)</i>	1	25	1	11.1	25
<b>Spionida</b>					
<b>Cirratulidae</b>					
<i>Caulleriella sp. J</i>	1	25	1	11.1	25
<b>Arthropoda</b>					
<b>Malacostraca</b>					
<b>Amphipoda</b>					
<b>Synopiidae</b>					
<i>Metatiron (LPIL)</i>	1	25	1	11.1	25
<b>Echinodermata</b>					
<b>Ophiuroidea</b>					
<i>Ophiuroidea (LPIL)</i>	1	25	1	11.1	25
<b>Mollusca</b>					
<b>Bivalvia</b>					
<b>Veneroida</b>					
<b>Lucinidae</b>					
<i>Lucina multilineata</i>	1	25	1	11.1	25
<b>Tellinidae</b>					
<i>Tellina (LPIL)</i>	1	25	1	11.1	25

**Station Data Summary Report****Station B2**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: B2  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Platyhelminthes					
Turbellaria					
Turbellaria (LPIL)					
	1	25	1	11.1	25
Rhynchocoela					
Rhynchocoela (LPIL)					
	1	25	1	11.1	25
Sipuncula					
Aspidosiphonidae					
Aspidosiphon albus					
	1	25	1	11.1	25

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station B2**

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 3

Station: B2  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	9	9	9	0
Total Individuals	9	9	9	0
Density (nos/sq.m.)		225	225	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' =	2.2
Species Diversity (Shannon; log base 2)	d =	3.17
Species Diversity (Shannon; log base 10)	H =	0.95
Species Diversity (Simpson; 1/S)	1/S =	
Species Evenness (Pielou)	J' =	1
Species Richness (Margalef)	D =	3.64
Equitability Index (Lloyd & Ghelardi)	e =	1.41

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
		% Total	Individuals	% Total
Platyhelminthes	1	11.1	1	11.1
Rhynchocoela	1	11.1	1	11.1
Annelida	2	22.2	2	22.2
Mollusca	2	22.2	2	22.2
Sipuncula	1	11.1	1	11.1
Arthropoda	1	11.1	1	11.1
Echinodermata	1	11.1	1	11.1
<b>TOTALS</b>	<b>9</b>		<b>9</b>	

### Station Data Summary Report

#### Station B3

Client: Wilmington District, USACE

Page 1

Station: B3

Project: Topsail Benthic Survey

Sample Type: Macrofauna

Location:

Replicates: 1

Sample Date: 28-29 Nov 06

Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Annelida					
Oligochaeta					
Tubificida					
Enchytraeidae					
Enchytraeidae (LPIL)	2	50	2	3.9	50
Tubificidae					
Tubificidae (LPIL)	1	25	1	2	25
Polychaeta					
Eunicida					
Onuphidae					
Onuphidae (LPIL)	1	25	1	2	25
Opheliida					
Opheliidae					
Armandia maculata	1	25	1	2	25
Orbiniida					
Paraonidae					
Aricidea catherinae	1	25	1	2	25
Cirrophorus (LPIL)	1	25	1	2	25
Phyllodocida					
Chrysopetalidae					
Bhawania heteroseta	5	125	5	9.8	125
Glyceridae					
Glyceridae (LPIL)	1	25	1	2	25
Goniadidae					
Goniadides caroliniae	7	175	7	13.7	175
Nereidae					
Nereididae (LPIL)	1	25	1	2	25

### Station Data Summary Report

#### Station B3

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: B3  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Phyllodocidae					
<i>Eumida sanguinea</i>	1	25	1	2	25
Pilargidae					
<i>Ancistrosyllis (LPIL)</i>	1	25	1	2	25
<i>Ancistrosyllis hartmanae</i>	1	25	1	2	25
Spionida					
Cirratulidae					
<i>Cirratulidae (LPIL)</i>	1	25	1	2	25
Spionidae					
<i>Spionidae (LPIL)</i>	4	100	4	7.8	100
Terebellida					
Terebellidae					
<i>Terebellidae (LPIL)</i>	1	25	1	2	25
Arthropoda					
Malacostraca					
Amphipoda					
Aoridae					
<i>Unciola serrata</i>	1	25	1	2	25
Isaeidae					
<i>Microprotopus raneyi</i>	2	50	2	3.9	50
Decapoda					
Paguridae					
<i>Paguridae (LPIL)</i>	1	25	1	2	25
Bryozoa					
<i>Bryozoa (LPIL)</i>	1	25	1	2	25

### Station Data Summary Report

#### Station B3

Client: Wilmington District, USACE

Page 1

Station: B3

Project: Topsail Benthic Survey

Sample Type: Macrofauna

Location: Topsail Beach

Replicates: 1

Sample Date: 28-29 Nov 06

Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Echinodermata					
Ophiuroidea					
Ophiuroidea (LPIL)	5	125	5	9.8	125
Mollusca					
Bivalvia					
Bivalvia (LPIL)	1	25	1	2	25
Veneroida					
Crassatellidae					
Crassinella lunulata	6	150	6	11.8	150
Semelidae					
Semelidae (LPIL)	1	25	1	2	25
Gastropoda					
Mesogastropoda					
Naticidae					
Tectonatica pusilla	1	25	1	2	25
Phoronida					
Phoronidae					
Phoronis (LPIL)	1	25	1	2	25
Platyhelminthes					
Turbellaria					
Turbellaria (LPIL)	1	25	1	2	25

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report**

**Station B3**

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 4

Station: B3  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	27	27	27	0
Total Individuals	51	51	51	0
Density (nos/sq.m.)		1275	1275	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' =	2.98
Species Diversity (Shannon; log base 2)	d =	4.29
Species Diversity (Shannon; log base 10)	H =	1.29
Species Diversity (Simpson; 1/S)	1/S =	19.92
Species Evenness (Pielou)	J' =	0.9
Species Richness (Margalef)	D =	6.61
Equitability Index (Lloyd & Ghelardi)	e =	1.06

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Platyhelminthes	1	3.7	1	1.9
Annelida	16	59.2	30	58.8
Mollusca	4	14.8	9	17.6
Arthropoda	3	11.1	4	7.8
Phoronida	1	3.7	1	1.9
Bryozoa	1	3.7	1	1.9
Echinodermata	1	3.7	5	9.8
<b>TOTALS</b>	<b>27</b>		<b>51</b>	

**Station Data Summary Report****Station C1**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: C1  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Annelida					
Polychaeta					
Phyllodocida					
Goniadidae					
<i>Goniada littorea</i>	2	50	2	40	50
Arthropoda					
Malacostraca					
Amphipoda					
Phoxocephalidae					
<i>Rhepoxynius hudsoni</i>	2	50	2	40	50
Mollusca					
Gastropoda					
Cephalaspidea					
Scaphandridae					
<i>Cyllichna alba</i>	1	25	1	20	25

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station C1**

Page 2

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Station: C1  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	3	3	3	0
Total Individuals	5	5	5	0
Density (nos/sq.m.)		125	125	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 1.05
Species Diversity (Shannon; log base 2)	d = 1.52
Species Diversity (Shannon; log base 10)	H = 0.46
Species Diversity (Simpson; 1/S)	1/S = 5
Species Evenness (Pielou)	J' = 0.96
Species Richness (Margalef)	D = 1.24
Equitability Index (Lloyd & Ghelardi)	e = 1.29

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Annelida	1	33.3	2	40
Mollusca	1	33.3	1	20
Arthropoda	1	33.3	2	40
<b>TOTALS</b>	<b>3</b>		<b>5</b>	

## Station Data Summary Report

### Station C2

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: C2  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Phyllodocida					
Goniadidae					
<i>Goniada littorea</i>	3	75	3	17.6	75
<b>Arthropoda</b>					
Malacostraca					
Amphipoda					
Amphipoda (LPIL)	1	25	1	5.9	25
Oedicerotidae					
Oedicerotidae (LPIL)	1	25	1	5.9	25
Platyischnopidae					
<i>Eudevenopus honduranus</i>	3	75	3	17.6	75
Decapoda					
Pasiphaeidae					
<i>Leptochela serratorbita</i>	1	25	1	5.9	25
Ostracoda					
Myodocopina					
Sarsiellidae					
<i>Eusarsiella texana</i>	1	25	1	5.9	25
<b>Echinodermata</b>					
Ophiuroidea					
Ophiuroidea (LPIL)	2	50	2	11.8	50

**Station Data Summary Report****Station C2**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: C2  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Annelida					
Mollusca					
Bivalvia					
Veneroida					
Tellinidae					
<i>Tellina</i> (LPIL)	4	100	4	23.5	100
Gastropoda					
Mesogastropoda					
Caecidae					
<i>Caecum pulchellum</i>	1	25	1	5.9	25

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station C2**

Page 3

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Station: C2  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	9	9	9	0
Total Individuals	17	17	17	0
Density (nos/sq.m.)		425	425	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 2.04
Species Diversity (Shannon; log base 2)	d = 2.94
Species Diversity (Shannon; log base 10)	H = 0.88
Species Diversity (Simpson; 1/S)	1/S = 10.46
Species Evenness (Pielou)	J' = 0.93
Species Richness (Margalef)	D = 2.82
Equitability Index (Lloyd & Ghelardi)	e = 1.2

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Annelida	1	11.1	3	17.6
Mollusca	2	22.2	5	29.4
Arthropoda	5	55.5	7	41.1
Echinodermata	1	11.1	2	11.7
<b>TOTALS</b>	<b>9</b>		<b>17</b>	

### Station Data Summary Report

#### Station C3

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: C3  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Eunicida					
Onuphidae					
Onuphidae (LPIL)	1	25	1	6.3	25
Phyllodocida					
Glyceridae					
Glyceridae (LPIL)	1	25	1	6.3	25
<b>Arthropoda</b>					
Malacostraca					
Amphipoda					
Platysischnopidae					
Eudevenopus honduranus	3	75	3	18.8	75
<b>Echinodermata</b>					
Echinoidea					
Echinoidea (LPIL)	2	50	2	12.5	50
Ophiuroidea					
Ophiuroidea (LPIL)	1	25	1	6.3	25
<b>Mollusca</b>					
Bivalvia					
Veneroida					
Tellinidae					
Tellina (LPIL)	5	125	5	31.3	125

**Station Data Summary Report****Station C3**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: C3  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Gastropoda					
Cephalaspidea					
Scaphandriidae					
Acteocina canaliculata	1	25	1	6.3	25
Mesogastropoda					
Caecidae					
Caecum pulchellum	1	25	1	6.3	25
Rhynchocoela					
Rhynchocoela (LPIL)	1	25	1	6.3	25

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station C3**

Page 3

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Station: C3  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	9	9	9	0
Total Individuals	16	16	16	0
Density (nos/sq.m.)		400	400	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 1.98
Species Diversity (Shannon; log base 2)	d = 2.85
Species Diversity (Shannon; log base 10)	H = 0.86
Species Diversity (Simpson; 1/S)	1/S = 8.57
Species Evenness (Pielou)	J' = 0.9
Species Richness (Margalef)	D = 2.89
Equitability Index (Lloyd & Ghelardi)	e = 1.12

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Rhynchocoela	1	11.1	1	6.2
Annelida	2	22.2	2	12.5
Mollusca	3	33.3	7	43.7
Arthropoda	1	11.1	3	18.7
Echinodermata	2	22.2	3	18.7
<b>TOTALS</b>	<b>9</b>		<b>16</b>	

### Station Data Summary Report

#### Station C4

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: C4  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Eunicida					
Onuphidae					
Onuphidae (LPIL)	1	25	1	5.9	25
Opheliida					
Opheliidae					
Armandia maculata	3	75	3	17.6	75
Spionida					
Chaetopteridae					
Spiochaetopterus oculatus	1	25	1	5.9	25
<b>Arthropoda</b>					
Malacostraca					
Amphipoda					
Platysischnopidae					
Eudevenopus honduranus	4	100	4	23.5	100
Decapoda					
Pinnotheridae					
Dissodactylus mellitae	2	50	2	11.8	50
Ostracoda					
Myodocopina					
Sarsiellidae					
Eusarsiella texana	1	25	1	5.9	25
Podocopida					
Podocopida (LPIL)	1	25	1	5.9	25

**Station Data Summary Report****Station C4**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: C4  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Echinodermata					
Echinoidea					
Clypeasteroida					
Mellitidae					
<i>Mellita isometra</i>	1	25	1	5.9	25
Mollusca					
Gastropoda					
Cephalaspidea					
Scaphandridae					
<i>Cylichna alba</i>	1	25	1	5.9	25
Mesogastropoda					
Caecidae					
<i>Caecum pulchellum</i>	2	50	2	11.8	50

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station C4**

Page 3

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Station: C4  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	10	10	10	0
Total Individuals	17	17	17	0
Density (nos/sq.m.)		425	425	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' =	2.15
Species Diversity (Shannon; log base 2)	d =	3.1
Species Diversity (Shannon; log base 10)	H =	0.93
Species Diversity (Simpson; 1/S)	1/S =	12.36
Species Evenness (Pielou)	J' =	0.93
Species Richness (Margalef)	D =	3.18
Equitability Index (Lloyd & Ghelardi)	e =	1.21

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Annelida	3	30	5	29.4
Mollusca	2	20	3	17.6
Arthropoda	4	40	8	47
Echinodermata	1	10	1	5.8
<b>TOTALS</b>	<b>10</b>		<b>17</b>	

**Station Data Summary Report****Station D1**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: D1  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Annelida					
Polychaeta					
Oweniida					
Oweniidae					
<i>Owenia fusiformis</i>	1	25	1	25	25
Arthropoda					
Malacostraca					
Amphipoda					
Platyischnopidae					
<i>Eudevenopus honduranus</i>	1	25	1	25	25
Ostracoda					
Myodocopina					
Sarsiellidae					
<i>Eusarsiella texana</i>	1	25	1	25	25
Mollusca					
Gastropoda					
Cephalaspidea					
Scaphandridae					
<i>Cylichna alba</i>	1	25	1	25	25

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

## Station Data Summary Report

### Station D1

Client: Wilmington District, USACE

Project: Topsail Benthic Survey

Location: Topsail Beach

Sample Date: 28-29 Nov 06

Page 2

Station: D1

Sample Type: Macrofauna

Replicates: 1

Sample Area: 0.0400

### **DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	4	4	4	0
Total Individuals	4	4	4	0
Density (nos/sq.m.)		100	100	0

### **FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 1.39
Species Diversity (Shannon; log base 2)	d = 2
Species Diversity (Shannon; log base 10)	H = 0.6
Species Diversity (Simpson; 1/S)	1/S =
Species Evenness (Pielou)	J' = 1
Species Richness (Margalef)	D = 2.16
Equitability Index (Lloyd & Ghelardi)	e = 1.37

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Annelida	1	25	1	25
Mollusca	1	25	1	25
Arthropoda	2	50	2	50
<b>TOTALS</b>	<b>4</b>		<b>4</b>	

## Station Data Summary Report

### Station D2

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: D2  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
<b>Polychaeta</b>					
<b>Eunicida</b>					
<b>Onuphidae</b>					
Onuphidae (LPIL)	1	25	1	10	25
<b>Phyllodocida</b>					
<b>Goniadidae</b>					
Goniada littorea	1	25	1	10	25
<b>Spionida</b>					
<b>Magelonidae</b>					
Magelona papillicornis	3	75	3	30	75
<b>Spiophanidae</b>					
Spiophanes bombyx	1	25	1	10	25
<b>Mollusca</b>					
<b>Bivalvia</b>					
<b>Veneroida</b>					
<b>Lucinidae</b>					
Lucinidae (LPIL)	1	25	1	10	25
<b>Mesodesmatidae</b>					
Ervilia concentrica	1	25	1	10	25
<b>Gastropoda</b>					
<b>Mesogastropoda</b>					
<b>Caecidae</b>					
Caecum pulchellum	1	25	1	10	25

**Station Data Summary Report****Station D2**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location:  
Sample Date: 28-29 Nov 06

Page 1

Station: D2  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Rhynchocoela					
Rhynchocoela (LPIL)	1	25	1	10	25

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

## Station Data Summary Report

### Station D2

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 3

Station: D2  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

#### **DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	8	8	8	0
Total Individuals	10	10	10	0
Density (nos/sq.m.)		250	250	0

#### **FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 1.97
Species Diversity (Shannon; log base 2)	d = 2.85
Species Diversity (Shannon; log base 10)	H = 0.86
Species Diversity (Simpson; 1/S)	1/S = 15
Species Evenness (Pielou)	J' = 0.95
Species Richness (Margalef)	D = 3.04
Equitability Index (Lloyd & Ghelardi)	e = 1.26

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Rhynchocoela	1	12.5	1	10
Annelida	4	50	6	60
Mollusca	3	37.5	3	30
<b>TOTALS</b>	<b>8</b>		<b>10</b>	

**Station Data Summary Report****Station D3**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: D3  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Annelida					
Polychaeta					
Phyllodocida					
Goniadidae					
<i>Goniada littorea</i>	2	50	2	28.6	50
Spionida					
Spionidae					
<i>Spionidae (LPIL)</i>	1	25	1	14.3	25
Mollusca					
Gastropoda					
Cephalaspidea					
Scaphandridae					
<i>Acteocina canaliculata</i>	1	25	1	14.3	25
Mesogastropoda					
Caecidae					
<i>Caecum pulchellum</i>	3	75	3	42.9	75

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station D3**

Page 2

Station: D3

Sample Type: Macrofauna

Replicates: 1

Sample Area: 0.0400

Client: Wilmington District, USACE

Project: Topsail Benthic Survey

Location: Topsail Beach

Sample Date: 28-29 Nov 06

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	4	4	4	0
Total Individuals	7	7	7	0
Density (nos/sq.m.)		175	175	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' =	1.28
Species Diversity (Shannon; log base 2)	d =	1.84
Species Diversity (Shannon; log base 10)	H =	0.55
Species Diversity (Simpson; 1/S)	1/S =	5.25
Species Evenness (Pielou)	J' =	0.92
Species Richness (Margalef)	D =	1.54
Equitability Index (Lloyd & Ghelardi)	e =	1.22

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No. Taxa</b>	<b>Taxa % Total</b>	<b>Total No. Individuals</b>	<b>Individuals % Total</b>
Annelida	2	50	3	42.8
Mollusca	2	50	4	57.1
<b>TOTALS</b>	<b>4</b>		<b>7</b>	

### Station Data Summary Report

#### Station D4

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: D4  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Eunicida					
Lumbrineridae					
<i>Lumbrineris latreilli</i>	1	25	1	5	25
Orbiniida					
Paraonidae					
<i>Aricidea (LPIL)</i>	1	25	1	5	25
<i>Aricidea wassi</i>	1	25	1	5	25
Phyllodocida					
Goniadidae					
<i>Goniada littorea</i>	1	25	1	5	25
Spionida					
Magelonidae					
<i>Magelona papillicornis</i>	1	25	1	5	25
<b>Arthropoda</b>					
Malacostraca					
Amphipoda					
<i>Amphipoda (LPIL)</i>	1	25	1	5	25
Oedicerotidae					
<i>Americhelidium americanum</i>	1	25	1	5	25
Phoxocephalidae					
<i>Rhepoxygnus hudsoni</i>	1	25	1	5	25
Platyischnopidae					
<i>Eudevenopus honduranus</i>	3	75	3	15	75

### Station Data Summary Report

#### Station D4

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: D4  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Cumacea					
Diastylidae					
Oxyurostylis (LPIL)	1	25	1	5	25
Ostracoda					
Myodocopina					
Philomedidae					
Harbansus paucichelatus	1	25	1	5	25
Philomedidae (LPIL)	1	25	1	5	25
Echinodermata					
Ophiuroidea					
Ophiuroidea (LPIL)	1	25	1	5	25
Mollusca					
Bivalvia					
Veneroida					
Lucinidae					
Lucina (LPIL)	1	25	1	5	25
Tellinidae					
Tellina (LPIL)	1	25	1	5	25
Gastropoda					
Mesogastropoda					
Caecidae					
Caecum pulchellum	3	75	3	15	75

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station D4**

Page 2

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Station: D4  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	16	16	16	0
Total Individuals	20	20	20	0
Density (nos/sq.m.)		500	500	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 2.67
Species Diversity (Shannon; log base 2)	d = 3.85
Species Diversity (Shannon; log base 10)	H = 1.16
Species Diversity (Simpson; 1/S)	1/S = 31.67
Species Evenness (Pielou)	J' = 0.96
Species Richness (Margalef)	D = 5.01
Equitability Index (Lloyd & Ghelardi)	e = 1.3

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Annelida	5	31.2	5	25
Mollusca	3	18.7	5	25
Arthropoda	7	43.7	9	45
Echinodermata	1	6.2	1	5
<b>TOTALS</b>	<b>16</b>		<b>20</b>	

## Station Data Summary Report

### Station E1

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: E1  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Orbiniida					
Paraonidae					
Aricidea wassi	1	25	1	6.7	25
Phyllodocida					
Glyceridae					
Glyceridae (LPIL)	1	25	1	6.7	25
Goniadidae					
Goniada littorea	3	75	3	20	75
Pilargidae					
Ancistrosyllis (PIL)	1	25	1	6.7	25
Spionida					
Magelonidae					
Magelona papillicornis	1	25	1	6.7	25
Spionidae					
Spionidae (LPIL)	1	25	1	6.7	25
<b>Arthropoda</b>					
Malacostraca					
Amphipoda					
Oedicerotidae					
Americhelidium americanum	1	25	1	6.7	25
Platyischnopidae					
Eudevenopus honduranus	1	25	1	6.7	25

**Station Data Summary Report****Station E1**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location:  
Sample Date: 28-29 Nov 06

Page 1

Station: E1  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Decapoda					
Paguridae					
Pagurus (LPIL)	1	25	1	6.7	25
Mollusca					
Bivalvia					
Veneroida					
Lucinidae					
Lucinidae (LPIL)	1	25	1	6.7	25
Gastropoda					
Mesogastropoda					
Caecidae					
Caecum pulchellum	3	75	3	20	75

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station E1**

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 3

Station: E1  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	11	11	11	0
Total Individuals	15	15	15	0
Density (nos/sq.m.)		375	375	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 2.27
Species Diversity (Shannon; log base 2)	d = 3.27
Species Diversity (Shannon; log base 10)	H = 0.99
Species Diversity (Simpson; 1/S)	1/S = 17.5
Species Evenness (Pielou)	J' = 0.95
Species Richness (Margalef)	D = 3.69
Equitability Index (Lloyd & Ghelardi)	e = 1.25

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Annelida	6	54.5	8	53.3
Mollusca	2	18.1	4	26.6
Arthropoda	3	27.2	3	20
<b>TOTALS</b>	<b>11</b>		<b>15</b>	

### Station Data Summary Report

#### Station E2

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: E2  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Capitellida					
Capitellidae					
<i>Mediomastus</i> (LPIL)	1	25	1	3.8	25
Opheliida					
Opheliidae					
<i>Opheliidae</i> (LPIL)	1	25	1	3.8	25
Orbiniida					
Paraonidae					
<i>Aricidea</i> (LPIL)	1	25	1	3.8	25
<i>Aricidea wassi</i>	2	50	2	7.7	50
Spionida					
Magelonidae					
<i>Magelona papillicornis</i>	2	50	2	7.7	50
<b>Arthropoda</b>					
Malacostraca					
Amphipoda					
Platysischnopidae					
<i>Eudevenopus honduranus</i>	3	75	3	11.5	75
<b>Echinodermata</b>					
Ophiuroidea					
<i>Ophiuroidea</i> (LPIL)	1	25	1	3.8	25

## Station Data Summary Report

### Station E2

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location:  
 Sample Date: 28-29 Nov 06

Page 1

Station: E2  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Mollusca</b>					
<b>Bivalvia</b>					
Veneroida					
Lucinidae					
Lucinidae (LPIL)	2	50	2	7.7	50
Tellinidae					
Tellina (LPIL)	1	25	1	3.8	25
Gastropoda					
Cephalaspidea					
Scaphandridae					
Acteocina canaliculata	2	50	2	7.7	50
Mesogastropoda					
Caecidae					
Caecum pulchellum	7	175	7	26.9	175
Eulimidae					
Strombiformis bilineatus	1	25	1	3.8	25
Rhynchocoela					
Rhynchocoela (LPIL)	2	50	2	7.7	50

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station E2**

Page 3

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Station: E2  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	13	13	13	0
Total Individuals	26	26	26	0
Density (nos/sq.m.)		650	650	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 2.34
Species Diversity (Shannon; log base 2)	d = 3.38
Species Diversity (Shannon; log base 10)	H = 1.02
Species Diversity (Simpson; 1/S)	1/S = 11.21
Species Evenness (Pielou)	J' = 0.91
Species Richness (Margalef)	D = 3.68
Equitability Index (Lloyd & Ghelardi)	e = 1.14

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Rhynchocoela	1	7.6	2	7.6
Annelida	5	38.4	7	26.9
Mollusca	5	38.4	13	50
Arthropoda	1	7.6	3	11.5
Echinodermata	1	7.6	1	3.8
<b>TOTALS</b>	<b>13</b>		<b>26</b>	

### Station Data Summary Report

#### Station E3

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: E3  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Capitellida					
Maldanidae					
Maldanidae (LPIL)	1	25	1	2.3	25
Opheliida					
Opheliidae					
Armandia (LPIL)	1	25	1	2.3	25
Orbiniida					
Paraonidae					
Aricidea suecica	1	25	1	2.3	25
Cirrophorus (LPIL)	1	25	1	2.3	25
Phyllodocida					
Goniadidae					
Goniada littorea	1	25	1	2.3	25
Pilargiidae					
Ancistrosyllis hartmanae	1	25	1	2.3	25
Spionida					
Magelonidae					
Magelona papillicornis	2	50	2	4.7	50
<b>Arthropoda</b>					
Malacostraca					
Amphipoda					
Phoxocephalidae					
Metharpinia floridana	1	25	1	2.3	25

### Station Data Summary Report

#### Station E3

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: E3  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Platyischnopidae					
Eudevenopus honduranus	6	150	6	14	150
Cumacea					
Diastylidae					
Oxyurostylis (LPIL)	1	25	1	2.3	25
Chordata					
Leptocardia					
Amphioxo					
Branchiostomidae					
Branchiostoma (LPIL)	2	50	2	4.7	50
Echinodermata					
Echinodermata (LPIL)	1	25	1	2.3	25
Mollusca					
Bivalvia					
Veneroida					
Crassatellidae					
Crassinella dupliniana	3	75	3	7	75
Crassinella lunulata	2	50	2	4.7	50
Montacutidae					
Montacutidae (LPIL)	1	25	1	2.3	25
Tellinidae					
Strigilla mirabilis	1	25	1	2.3	25
Tellina (LPIL)	1	25	1	2.3	25

**Station Data Summary Report****Station E3**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: E3  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Gastropoda					
Cephalaspidea					
Scaphandridae					
<i>Acteocina canaliculata</i>	1	25	1	2.3	25
Mesogastropoda					
Caecidae					
<i>Caecum pulchellum</i>	12	300	12	27.9	300
Eulimidae					
<i>Strombiformis bilineatus</i>	1	25	1	2.3	25
Naticidae					
<i>Tectonatica pusilla</i>	1	25	1	2.3	25
Rhynchocoela					
<i>Rhynchocoela (LPIL)</i>	1	25	1	2.3	25

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station E3**

Page 4

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Station: E3  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	22	22	22	0
Total Individuals	43	43	43	0
Density (nos/sq.m.)		1075	1075	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' =	2.64
Species Diversity (Shannon; log base 2)	d =	3.82
Species Diversity (Shannon; log base 10)	H =	1.15
Species Diversity (Simpson; 1/S)	1/S =	10.38
Species Evenness (Pielou)	J' =	0.86
Species Richness (Margalef)	D =	5.58
Equitability Index (Lloyd & Ghelardi)	e =	0.92

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No. Taxa</b>	<b>Taxa % Total</b>	<b>Total No. Individuals</b>	<b>Individuals % Total</b>
Rhynchocoela	1	4.5	1	2.3
Annelida	7	31.8	8	18.6
Mollusca	9	40.9	23	53.4
Arthropoda	3	13.6	8	18.6
Echinodermata	1	4.5	1	2.3
Chordata	1	4.5	2	4.6
<b>TOTALS</b>	<b>22</b>		<b>43</b>	

## Station Data Summary Report

### Station E4

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: E4  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Eunicida					
Oenonidae					
<i>Arabella multidentata</i>	1	25	1	3.6	25
Opheliida					
Opheliidae					
<i>Armandia maculata</i>	2	50	2	7.1	50
<i>Opheliidae (LPIL)</i>	1	25	1	3.6	25
Phyllodocida					
Nereidae					
<i>Ceratocephale oculata</i>	1	25	1	3.6	25
Spionida					
Magelonidae					
<i>Magelona papillicornis</i>	2	50	2	7.1	50
Spionidae					
<i>Parapriionospio pinnata</i>	1	25	1	3.6	25
<b>Arthropoda</b>					
Malacostraca					
Amphipoda					
Phoxocephalidae					
<i>Rhepoxynius hudsoni</i>	1	25	1	3.6	25
Platyischnopidae					
<i>Eudevenopus honduranus</i>	7	175	7	25	175

## Station Data Summary Report

### Station E4

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: E4  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Ostracoda					
Myodocopina					
Sarsiellidae					
<i>Eusarsiella texana</i>	1	25	1	3.6	25
Mollusca					
Bivalvia					
Veneroida					
Crassatellidae					
<i>Crassinella dupliniana</i>	1	25	1	3.6	25
Lucinidae					
<i>Lucina multilineata</i>	3	75	3	10.7	75
Tellinidae					
<i>Strigilla mirabilis</i>	1	25	1	3.6	25
<i>Tellina (LPIL)</i>	1	25	1	3.6	25
Gastropoda					
Cephalaspidea					
Scaphandridae					
<i>Acteocina canaliculata</i>	1	25	1	3.6	25
Mesogastropoda					
Caecidae					
<i>Caecum pulchellum</i>	2	50	2	7.1	50
Naticidae					
<i>Tectonatica pusilla</i>	1	25	1	3.6	25

**Station Data Summary Report****Station E4**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: E4  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Rhynchocoela					
Rhynchocoela (LPIL)	1	25	1	3.6	25

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station E4**

Page 4

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Station: E4  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	17	17	17	0
Total Individuals	28	28	28	0
Density (nos/sq.m.)		700	700	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' =	2.58
Species Diversity (Shannon; log base 2)	d =	3.72
Species Diversity (Shannon; log base 10)	H =	1.12
Species Diversity (Simpson; 1/S)	1/S =	14
Species Evenness (Pielou)	J' =	0.91
Species Richness (Margalef)	D =	4.8
Equitability Index (Lloyd & Ghelardi)	e =	1.12

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Rhynchoela	1	5.8	1	3.5
Annelida	6	35.2	8	28.5
Mollusca	7	41.1	10	35.7
Arthropoda	3	17.6	9	32.1
<b>TOTALS</b>	<b>17</b>		<b>28</b>	

### Station Data Summary Report

#### Station F1

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: F1  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Opheliida					
Opheliidae					
Opheliidae (LPIL)	1	25	1	11.1	25
Spionida					
Cirratulidae					
Caullerella sp. J	1	25	1	11.1	25
Magelonidae					
Magelona papillicornis	1	25	1	11.1	25
Magelona pettiboneae	1	25	1	11.1	25
<b>Arthropoda</b>					
Malacostraca					
Cumacea					
Diastylidae					
Oxyurostylis (LPIL)	1	25	1	11.1	25
Ostracoda					
Myodocopina					
Sarsiellidae					
Eusarsiella texana	1	25	1	11.1	25
<b>Chordata</b>					
Leptocardia					
Amphioxo					
Branchiostomidae					
Branchiostoma (LPIL)	1	25	1	11.1	25

**Station Data Summary Report****Station F1**

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: F1  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

<b>TAXON</b>	<b>Rep 1</b>		<b>Station</b>		
	Count	Density	Total	Percent	Mean Density
Mollusca					
Bivalvia					
Veneroida					
Crassatellidae					
<i>Crassinella lunulata</i>	1	25	1	11.1	25
Platyhelminthes					
Turbellaria					
<i>Turbellaria (LPIL)</i>	1	25	1	11.1	25

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station F1**

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 3

Station: F1  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	9	9	9	0
Total Individuals	9	9	9	0
Density (nos/sq.m.)		225	225	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 2.2
Species Diversity (Shannon; log base 2)	d = 3.17
Species Diversity (Shannon; log base 10)	H = 0.95
Species Diversity (Simpson; 1/S)	1/S =
Species Evenness (Pielou)	J' = 1
Species Richness (Margalef)	D = 3.64
Equitability Index (Lloyd & Ghelardi)	e = 1.41

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Platyhelminthes	1	11.1	1	11.1
Annelida	4	44.4	4	44.4
Mollusca	1	11.1	1	11.1
Arthropoda	2	22.2	2	22.2
Chordata	1	11.1	1	11.1
<b>TOTALS</b>	<b>9</b>		<b>9</b>	

Station Data Summary Report						
Station F2			Station F2			
Client: Wilmington District, USACE		Page 1		Station: F2		
Project: Topsail Benthic Survey		Sample Type: Macrofauna		Replicates: 1		
Location: Topsail Beach		Sample Area: 0.0400				
Sample Date: 28-29 Nov 06						
TAXON	Rep 1		Station			
	Count	Density	Total	Percent	Mean Density	
Annelida						
Polychaeta						
Opheliida						
Opheliidae						
<i>Ophelia denticulata</i>	1	25	1	4.2	25	
Phyllodocida						
Chrysopetalidae						
<i>Bhawania heteroseta</i>	1	25	1	4.2	25	
Goniadidae						
<i>Goniadides carolinae</i>	2	50	2	8.3	50	
Hesionidae						
<i>Microphthalmus (LPIL)</i>	1	25	1	4.2	25	
Nephtyidae						
<i>Nephtys simoni</i>	1	25	1	4.2	25	
Spionida						
Cirratulidae						
<i>Cirratulidae (LPIL)</i>	1	25	1	4.2	25	
Arthropoda						
Malacostraca						
Amphipoda						
Melitidae						
<i>Maera caroliniana</i>	1	25	1	4.2	25	

**Station Data Summary Report****Station F2**

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 4

Station: F2  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	16	16	16	0
Total Individuals	24	24	24	0
Density (nos/sq.m.)		600	600	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 2.64
Species Diversity (Shannon; log base 2)	d = 3.8
Species Diversity (Shannon; log base 10)	H = 1.14
Species Diversity (Simpson; 1/S)	1/S = 23
Species Evenness (Pielou)	J' = 0.95
Species Richness (Margalef)	D = 4.72
Equitability Index (Lloyd & Ghelardi)	e = 1.26

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Rhynchocoela	1	6.2	3	12.5
Annelida	6	37.5	7	29.1
Mollusca	7	43.7	11	45.8
Arthropoda	1	6.2	1	4.1
Chordata	1	6.2	2	8.3
<b>TOTALS</b>	<b>16</b>		<b>24</b>	

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

## Station Data Summary Report

## Station F2

Page 1

Station: F2

Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Chordata					
Leptocardia					
Amphioxo					
Branchiostomidae					
Branchiostoma (LPIL)	2	50	2	8.3	50
Mollusca					
Bivalvia					
Bivalvia (LPIL)	1	25	1	4.2	25
Ostreoida					
Anomiidae					
Anomia simplex	1	25	1	4.2	25
Veneroida					
Crassatellidae					
Crassinella dupliniana	4	100	4	16.7	100
Crassinella lunulata	2	50	2	8.3	50
Gastropoda					
Cephalaspidea					
Scaphandridae					
Acteocina canaliculata	1	25	1	4.2	25
Mesogastropoda					
Caecidae					
Caecum pulchellum	1	25	1	4.2	25
Pyramidelloida					
Pyramidellidae					
Turbonilla (LPIL)	1	25	1	4.2	25

Station Data Summary Report

Station F2

Client: Wilmington District, USACE

Page 1

Station: F2

Project: Topsail Benthic Survey

Sample Type: Macrofauna

Location: Topsail Beach

Replicates: 1

Sample Date: 28-29 Nov 06

Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Rhynchocoela					
Rhynchocoela (LPIL)	3	75	3	12.5	75

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station F2**

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 4

Station: F2  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	16	16	16	0
Total Individuals	24	24	24	0
Density (nos/sq.m.)		600	600	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 2.64
Species Diversity (Shannon; log base 2)	d = 3.8
Species Diversity (Shannon; log base 10)	H = 1.14
Species Diversity (Simpson; 1/S)	1/S = 23
Species Evenness (Pielou)	J' = 0.95
Species Richness (Margalef)	D = 4.72
Equitability Index (Lloyd & Ghelardi)	e = 1.26

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Rhynchocoela	1	6.2	3	12.5
Annelida	6	37.5	7	29.1
Mollusca	7	43.7	11	45.8
Arthropoda	1	6.2	1	4.1
Chordata	1	6.2	2	8.3
<b>TOTALS</b>	<b>16</b>		<b>24</b>	

## Station Data Summary Report

### Station F3

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: F3  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
<b>Annelida</b>					
Polychaeta					
Eunicida					
Onuphidae					
Onuphidae (LPIL)	1	25	1	7.1	25
Phyllodocida					
Goniadidae					
Goniada littorea	1	25	1	7.1	25
Nereidae					
Ceratocephale oculata	1	25	1	7.1	25
Spionida					
Spionidae					
Parapriionospio pinnata	1	25	1	7.1	25
<b>Arthropoda</b>					
Malacostraca					
Amphipoda					
Platyischnopidae					
Eudevenopus honduranus	2	50	2	14.3	50
<b>Echinodermata</b>					
Ophiuroidea					
Ophiuroidea (LPIL)	2	50	2	14.3	50

## Station Data Summary Report

### Station F3

Client: Wilmington District, USACE  
Project: Topsail Benthic Survey  
Location: Topsail Beach  
Sample Date: 28-29 Nov 06

Page 1

Station: F3  
Sample Type: Macrofauna  
Replicates: 1  
Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Mollusca					
Bivalvia					
Veneroida					
Lucinidae					
Lucinidae (LPIL)	2	50	2	14.3	50
Gastropoda					
Mesogastropoda					
Caecidae					
Caecum pulchellum	2	50	2	14.3	50
Neogastropoda					
Olividae					
Olivella dealbata	2	50	2	14.3	50

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station F3**

Page 3

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Station: F3  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	9	9	9	0
Total Individuals	14	14	14	0
Density (nos/sq.m.)		350	350	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 2.14
Species Diversity (Shannon; log base 2)	d = 3.09
Species Diversity (Shannon; log base 10)	H = 0.93
Species Diversity (Simpson; 1/S)	1/S = 18.2
Species Evenness (Pielou)	J' = 0.98
Species Richness (Margalef)	D = 3.03
Equitability Index (Lloyd & Ghelardi)	e = 1.34

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Annelida	4	44.4	4	28.5
Mollusca	3	33.3	6	42.8
Arthropoda	1	11.1	2	14.2
Echinodermata	1	11.1	2	14.2
<b>TOTALS</b>	<b>9</b>		<b>14</b>	

### Station Data Summary Report

#### Station F4

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: F4  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Annelida					
Polychaeta					
Eunicida					
Lumbrineridae					
<i>Lumbrineris latreilli</i>	1	25	1	3.4	25
Oenonidae					
<i>Arabella multidentata</i>	1	25	1	3.4	25
Phyllodocida					
Goniadidae					
<i>Goniada littorea</i>	1	25	1	3.4	25
Nephtyidae					
<i>Nephtys picta</i>	2	50	2	6.9	50
Nereidae					
<i>Ceratocephale oculata</i>	1	25	1	3.4	25
Spionida					
Magelonidae					
<i>Magelona papillicornis</i>	1	25	1	3.4	25
Spionidae					
<i>Spionidae (LPIL)</i>	1	25	1	3.4	25
Arthropoda					
Malacostraca					
Amphipoda					
Haustoriidae					
<i>Acanthohaustorius intermedius</i>	2	50	2	6.9	50

### Station Data Summary Report

#### Station F4

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 1

Station: F4  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

TAXON	Rep 1		Station		
	Count	Density	Total	Percent	Mean Density
Phoxocephalidae					
<i>Rhepoxygnus hudsoni</i>	1	25	1	3.4	25
Platyischnopidae					
<i>Eudevenopus honduranus</i>	12	300	12	41.4	300
Ostracoda					
Myodocopina					
Sarsiellidae					
<i>Eusarsiella radiicosta</i>	1	25	1	3.4	25
Echinodermata					
Echinoidea					
Clypeasteroida					
Mellitidae					
<i>Mellitidae (LPIL)</i>	2	50	2	6.9	50
Mollusca					
Bivalvia					
Veneroida					
Lucinidae					
<i>Lucinidae (LPIL)</i>	1	25	1	3.4	25
Mesodesmatidae					
<i>Ervilia concentrica</i>	1	25	1	3.4	25
Gastropoda					
Neogastropoda					
Turridae					
<i>Kurtziella atrostyla</i>	1	25	1	3.4	25

Note: LPIL designates the LOWEST PRACTICAL IDENTIFICATION LEVEL

**Station Data Summary Report****Station F4**

Client: Wilmington District, USACE  
 Project: Topsail Benthic Survey  
 Location: Topsail Beach  
 Sample Date: 28-29 Nov 06

Page 3

Station: F4  
 Sample Type: Macrofauna  
 Replicates: 1  
 Sample Area: 0.0400

**DATA SUMMARY**

<b>FAUNAL PARAMETERS</b>	<b>Station</b>	<b>Rep 1</b>	<b>Mean</b>	<b>Std Dev</b>
Total Taxa	15	15	15	0
Total Individuals	29	29	29	0
Density (nos/sq.m.)		725	725	0

**FAUNAL INDICES**

Species Diversity (Shannon; log base e)	H' = 2.2
Species Diversity (Shannon; log base 2)	d = 3.17
Species Diversity (Shannon; log base 10)	H = 0.95
Species Diversity (Simpson; 1/S)	1/S = 5.88
Species Evenness (Pielou)	J' = 0.81
Species Richness (Margalef)	D = 4.16
Equitability Index (Lloyd & Ghelardi)	e = 0.85

<b>MAJOR TAXONOMIC GROUPS</b>	<b>Total No.</b>	<b>Taxa</b>	<b>Total No.</b>	<b>Individuals</b>
	<b>Taxa</b>	<b>% Total</b>	<b>Individuals</b>	<b>% Total</b>
Annelida	7	46.6	8	27.5
Mollusca	3	20	3	10.3
Arthropoda	4	26.6	16	55.1
Echinodermata	1	6.6	2	6.8
<b>TOTALS</b>	<b>15</b>		<b>29</b>	

**APPENDIX D**

**Supporting Data Tables**

**Table D-1. Species abundance and density at sampling stations within borrow sites A, B, and C off Topsail Beach, NC.**

Species List	Mean A										Mean B										Mean C														
	A1		A2		A3		A4		A5		B1		B2		B3		C1		C2		C3		C4												
	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density							
<b>ANNELEIDA</b>																																			
CLASS OLIGOCHAETA																																			
Order TUBIFICIDA																																			
FAMILY ENCHYTRAEIDAE																																			
<i>Enchytraeidae (LPIL)</i>																	2	50	.67	16.67															
FAMILY TUBIFICIDAE																																			
<i>Tubificidae (LPIL)</i>																	1	25	.33	8.33															
CLASS POLYCHAETA																																			
Order CAPITELLIDA																																			
FAMILY CAPITELLIDAE																																			
<i>Capitellidae (LPIL)</i>	1	25									.20	5.00																							
<i>Mediomastus (LPIL)</i>																	3	75	1	25															
<i>Mediomastus californiensis</i>											1	25					.20	5																	
FAMILY MALDANIDAE																																			
<i>Maldanidae (LPIL)</i>																																			
Order EUNICIDA																																			
FAMILY DORVILLEIDAE																																			
<i>Schistomerings pectinata</i>																																			
FAMILY LUMBRINERIDAE																																			
<i>Lumbrineris latreilli</i>																																			
FAMILY OENONIDAE																																			
<i>Arabella multidentata</i>																																			
FAMILY ONUPHIDAE																																			
<i>Onuphidae (LPIL)</i>											1	25					.20	5.00																	
Order OPHELIIDA																																			
FAMILY OPHELIIDAE																																			
<i>Opheliidae (LPIL)</i>																																			
<i>Armandia (LPIL)</i>																																			
<i>Armandia maculata</i>	1	25									.20	5.00																							
<i>Ophelia denticulata</i>																																			
Order ORBINIIDAE																																			
FAMILY PARAONIDIADAE																																			
<i>Aricidea (LPIL)</i>																																			
<i>Aricidea catherinae</i>											1	25	.20	5.00																					
<i>Aricidea suecica</i>																																			
<i>Aricidea wassi</i>																																			
<i>Cirrophorus (LPIL)</i>																																			

**Table D-1. (continued)**

Species List	Mean A										Mean B						Mean C															
	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density								
Order OWENIIDAE																																
FAMILY OWENIIDAE																																
<i>Owenia fusiformis</i>	1	25							.20	5.00																						
Order PHYLLODOCIDA																																
FAMILY CHRYSOPETALIDAE																																
<i>Bhawania heteroseta</i>											2	50			5	125	2.33	58.33														
FAMILY GLYCERIDAE																																
<i>Glyceridae (LPIL)</i>			2	50					.40	10.00					1	25	.33	8.33				1	25		.25	6.25						
FAMILY GONIADIDAE																																
<i>Goniada littorea</i>							1	25			.20	5.00									2	50	3	75		1.25	31.25					
<i>Goniadides carolinae</i>															2	50			7	175	3	75.00										
FAMILY HESIONIDAE																																
<i>Microphthalmus (LPIL)</i>																																
FAMILY NEPHTYIDAE																																
<i>Aglaophamus verrilli</i>			1	25					.20	5.00																						
<i>Nephrys picta</i>																																
<i>Nephys simoni</i>																																
FAMILY NEREIDAE																																
<i>Nereididae (LPIL)</i>																			1	25	.33	8.33										
<i>Ceratocephale oculata</i>																																
FAMILY PHYLLODOCIDAE																																
<i>Eumida sanguinea</i>																			1	25	.33	8.33										
FAMILY PILARGIIDAE																																
<i>Ancistrosyllis (LPIL)</i>																			1	25	.33	8.33										
<i>Ancistrosyllis hartmanae</i>																			1	25	.33	8.33										
<i>Sigambra tentaculata</i>								1	25	.20	5.00																					
Order SPIONIDA																																
FAMILY CHAETOPTERIDAE																																
<i>Spiochaetopterus oculus</i>	1	25							.20	5.00															1	25	.25	6.25				
FAMILY CIRRATULIDAE																																
<i>Cirratulidae (LPIL)</i>																1	25		1	25	.67	16.67										
<i>Caulieriella sp. J</i>	1	25	2	50		1	25		.80	20.00			1	25		1	25		.33	8.33												
FAMILY MAGELONIDAE																																
<i>Magelona (LPIL)</i>	2	50	1	25					.60	15.00																						
<i>Magelona papillicornis</i>						2	50		3	75	1.00	25.00																				
<i>Magelona pectiniferae</i>																																
FAMILY SPIONIDAE																																
<i>Spionidae (LPIL)</i>						1	25			.20	5.00						4	100	1.33	33.33												
<i>Parapriionospio pirnata</i>						1	25			.20	5.00																					
<i>Spiophanes bombyx</i>																																

**Table D-1.** (continued)

**Table D-1. (continued)**

Species List	A1	A2	A3	A4	A5	Mean A		B1	B2	B3	Mean B		C1	C2	C3	C4	Mean C	
	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density								
CLASS OSTRACODA																		
Order MYODOCOPINA																		
FAMILY PHIOMEDIDAE																		
<i>Philomedidae (LPIL)</i>																		
<i>Harbansus paucichelatus</i>																		
FAMILY SARSIELLIDAE																		
<i>Eusarsiella (LPIL)</i>	1	.25					.20	5.00										
<i>Eusarsiella radicosta</i>							1	.25	.20	5.00							1	.25
<i>Eusarsiella texana</i>																	1	.25
Order PODOCOPIDA																		
<i>Podocopida (LPIL)</i>																	1	.25
<b>BRYOZOA</b>																		
<i>Bryozoa (LPIL)</i>																		
<b>CHORDATA</b>																		
CLASS LEPTOCARDIA																		
Order AMPHIOXI																		
FAMILY BRANCHIOSTOMIDAE																		
<i>Branchiostoma (LPIL)</i>																		
<b>CNIDARIA</b>																		
CLASS HYDROZOA																		
<i>Hydrozoa (LPIL)</i>	1	.25					.20	5.00										
<b>ECHINODERMATA</b>																		
<i>Echinodermata (LPIL)</i>																		
CLASS ECHINOIDEA																		
<i>Echinoidea (LPIL)</i>																	2	.50
Order CLYPEASTEROIDA																		
FAMILY MELLITIDAE																		
<i>Mellitidae (LPIL)</i>																		
<i>Mellita isometra</i>																	1	.25
CLASS OPHIUROIDEA																		
<i>Ophiuroidea (LPIL)</i>																	2	.50
Order OPHIURIDA																		
FAMILY AMPHIURIDAE																		
<i>Amphuridae (LPIL)</i>																		
<b>MOLLUSCA</b>																		
CLASS BIVALVIA																		
<i>Bivalvia (LPIL)</i>	1	.25			1	.25		.40	10.00			1	.25	.33	8.33			
Order OSTREOIDA																		
FAMILY ANOMIIDAE																		
<i>Anomia simplex</i>																		

**Table D-1. (continued)**

Species List	A1	A2	A3	A4	A5	Mean A		B1	B2	B3	Mean B		C1	C2	C3	C4	Mean C		
	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density									
Order VENEROIDA																			
FAMILY CRASSATELLIDAE																			
<i>Crassinella dupliniana</i>																			
<i>Crassinella lunulata</i>									4	100		6	150	3.33	83.33				
FAMILY LUCINIDAE																			
<i>Lucinidae (LPIL)</i>	2	.50				.40	10.00												
<i>Lucina (LPIL)</i>																			
<i>Lucina multilineata</i>											1	25		.33	8.33				
FAMILY MESODESMATIDAE																			
<i>Ervilia concentrica</i>				1	25		.20	5.00											
FAMILY MONTACUTIDAE																			
<i>Montacutidae (LPIL)</i>																			
FAMILY SEMELIDAE																			
<i>Semelidae (LPIL)</i>																			
FAMILY TELLINIDAE																			
<i>Strigilla mirabilis</i>																			
<i>Tellina (LPIL)</i>	2	.50				3	75	1.0	25.00		1	25		.33	8.33			2.25	56.25
<i>Tellina iris</i>	1	.25					.20	5.00											
CLASS GASTROPODA																			
Order CEPHALASPIDEA																			
FAMILY SCAPHANDRIDAE																			
<i>Acteoцина canaliculata</i>			1	25			.20	5.00									1	.25	
<i>Cylicha alba</i>	4	100	1	25			1.0	25.00									1	.25	
Order MESOGASTROPODA																			
FAMILY CAECIDAE																			
<i>Caecum pulchellum</i>										1	25			.33	8.33		1	.25	
FAMILY EULIMIDAE																	1	.25	
<i>Strombiformis bilineatus</i>																	2	50	
FAMILY NATICIDAE																	1	100	
<i>Tectonatica pusilla</i>																		25.00	
Order NEOGASTROPODA																			
FAMILY OLIVIDAE																			
<i>Oliva sayana</i>				2	.50	1	75	.60	25.00										
<i>Olivella dealbata</i>																			
FAMILY TURRIDAE																			
<i>Kurtziella atrostyla</i>																			
Order PYRAMIDELLOIDA																			
FAMILY PYRAMIDELLIDAE																			
<i>Odostomia weberi</i>							1	.25	.20	5.00									
<i>Turbonilla (LPIL)</i>																			

**Table D-1. (concluded)**

Species List	Mean A										Mean B										Mean C										
	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	
<b>PHORONIDA</b>																															
FAMILY PHORONIDAE																															
<i>Phoronis (LPIL)</i>	1	25					.20	5.00					1	25	.33	8.33															
<b>PLATYHELMINTHES</b>																															
CLASS TURBELLARIA																															
<i>Turbellaria (LPIL)</i>									2	50	1	25	1	25	1.33	33.33															
<b>RHYNCHOCOELA</b>																															
<i>Rhynchocoela (LPIL)</i>	1	25					.20	5.00	2	50	1	25					1.00	25.00								1	25		.25	6.25	
<b>SIPUNCULA</b>																															
FAMILY ASPIDOSIPHONIDAE																															
<i>Aspidosiphon albus</i>																															

**Table D-2. Species abundance and density at sampling stations within borrow sites D, E, and F off Topsail Beach, NC.**

Species List	D1		D2		D3		D4		Mean D		E1		E2		E3		E4		Mean E		F1		F2		F3		F4		Mean F	
	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density		
<b>ANNELEIDA</b>																														
CLASS OLIGOCHAETA																														
Order TUBIFICIDA																														
FAMILY ENCHYTRAIEIDAE																														
<i>Enchytraeidae (LPIL)</i>																														
FAMILY TUBIFICIDAE																														
<i>Tubificidae (LPIL)</i>																														
CLASS POLYCHAETA																														
Order CAPITELLIDA																														
FAMILY CAPITELLIDAE																														
<i>Capitellidae (LPIL)</i>																														
<i>Mediomastus (LPIL)</i>																	1	25				.25	6.25							
<i>Mediomastus californiensis</i>																														
FAMILY MALDANIDAE																														
<i>Maldanidae (LPIL)</i>																														
Order EUNICIDA																														
FAMILY DORVILLEIDAE																														
<i>Schistomerings pectinata</i>																														
FAMILY LUMBRINERIDAE																														
<i>Lumbrineris latreilli</i>																	1	25	.25	6.25										
FAMILY OENONIDAE																														
<i>Arabella multidentata</i>																														
FAMILY ONUPHIDAE																														
<i>Onuphidae (LPIL)</i>	1	25																												
Order OPHELIIDA																														
FAMILY OPHELIIDAE																														
<i>Opheliidae (LPIL)</i>																	1	25												
<i>Armandia (LPIL)</i>																		1	25											
<i>Armandia maculata</i>																			2	50	.50	12.50								
<i>Ophelia denticulata</i>																														
Order ORBINIIDA																														
FAMILY PARAONIDAE																														
<i>Aricidea (LPIL)</i>																	1	25	.25	6.25										
<i>Aricidea catherinae</i>																														
<i>Aricidea suecica</i>																		1	25											
<i>Aricidea wassi</i>																	1	25	.25	6.25										
<i>Cirrophorus (LPIL)</i>																			1	25										
Order OWENIIDA																														
FAMILY OWENIIDAE																														
<i>Owenia fusiformis</i>	1	25																												

**Table D-2. (continued)**

Species List	D1				D2				D3				D4				Mean D		E1				E2				E3				E4				Mean E		F1				F2				F3				F4				Mean F	
	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density																												
Order PHYLLODOCIDA																																																						
FAMILY CHRYSOPETALIDAE																																																						
<i>Bhawania heteroseta</i>																																																						
FAMILY GLYCERIDAE																																																						
<i>Glyceridae (LPIL)</i>																																																						
FAMILY GONIADIDAE																																																						
<i>Goniada littorea</i>	1	.25	2	.50	1	.25	1.00	25.00	3	.75																																												
<i>Goniadides carolinae</i>																																																						
FAMILY HESIONIDAE																																																						
<i>Microphthalmus (LPIL)</i>																																																						
FAMILY NEPTYDIAE																																																						
<i>Aglaophamus verrilli</i>																																																						
<i>Nephlys picta</i>																																																						
<i>Nephlys simoni</i>																																																						
FAMILY NEREIDAE																																																						
<i>Nereididae (LPIL)</i>																																																						
<i>Ceratocephale oculata</i>																																																						
FAMILY PHYLLODOCIDAE																																																						
<i>Eumida sanguinea</i>																																																						
FAMILY PILARGIIDAE																																																						
<i>Ancistrosyllis (LPIL)</i>																																																						
<i>Ancistrosyllis hartmanae</i>																																																						
<i>Sigambla tentaculata</i>																																																						
Order SPIONIDA																																																						
FAMILY CHAETOPTERIDAE																																																						
<i>Spiochaetopterus oculatus</i>																																																						
FAMILY CIRRATULIDAE																																																						
<i>Cirratulidae (LPIL)</i>																																																						
<i>Caulteriella sp. J</i>																																																						
FAMILY MAGELONIDAE																																																						
<i>Magelona (LPIL)</i>																																																						
<i>Magelona papillicornis</i>	3	.75					1	.25	1.00	25.00	1	.25	2	.50	2	.50	2	.50	1.75	43.75	1	.25																																
<i>Magelona pettiboneae</i>																																																						
FAMILY SPIONIDAE																																																						
<i>Spionidae (LPIL)</i>																																																						
<i>Parapriionospio pinnata</i>																																																						
<i>Spiophanes bombyx</i>	1	.25																																																				
Order TEREBELLIDA																																																						
FAMILY PECTINARIIDAE																																																						
<i>Pectinaria gouldii</i>																																																						
FAMILY TEREBELLIDAE																																																						
<i>Terebellidae (LPIL)</i>																																																						

**Table D-2. (continued)**

Species List	D1		D2		D3		D4		Mean D		E1		E2		E3		E4		Mean E		F1		F2		F3		F4		Mean F			
	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density				
ARTHROPODA																																
CLASS MALACOSTRACA																																
Order AMPHIPODA																																
Amphipoda (LPIL)									1	.25	.25	6.25																				
FAMILY AORIDAE																																
<i>Rildardanus laminosa</i>																																
<i>Unciola serrata</i>																																
FAMILY HAUSTORIIDAE																																
<i>Acanthohaustorius intermedius</i>																																
FAMILY ISAEIDAE																																
<i>Microprotopus raneyi</i>																																
FAMILY MELITIDAE																																
<i>Maera caroliniana</i>																																
FAMILY OEDICEROTIDAE																																
<i>Oedicerotidae (LPIL)</i>																																
<i>Americhelidium americanum</i>									1	.25	.25	6.25	1	.25																		
FAMILY PHOXOCEPHALIDAE																																
<i>Metharpinia floridana</i>																																
<i>Rhepoxyinius hudsoni</i>									1	.25	.25	6.25																				
FAMILY PLATYISCHNOPIDAE																																
<i>Eudevenopus honduranus</i>	1	.25							3	.75	.75	18.75	1	.25	3	.75	6	150	7	175	4.25	106.25					2	50	12	300	3.50	87.50
FAMILY SYNOPIIDAE																																
<i>Metatiron (LPIL)</i>																																
<i>Metatiron triocellatus</i>																																
Order CUMACEA																																
FAMILY DIASTYLIDAE																																
<i>Oxyurostylis (LPIL)</i>									1	.25	.25	6.25																				
Order DECAPODA																																
FAMILY PAGURIDAE																																
<i>Pagurus (LPIL)</i>																																
FAMILY PASIPHAEIDAE																																
<i>Leptochela serratiorbita</i>																																
FAMILY PINNOTHERIDAE																																
<i>Dissodactylus mellitae</i>																																
CLASS OSTRACODA																																
Order MYODOCOPINA																																
FAMILY PHILOMEDIDAE																																
<i>Philomedidae (LPIL)</i>									1	.25	.25	6.25																				
<i>Harbansus paucichelatus</i>									1	.25	.25	6.25																				
FAMILY SARSIELLIIDAE																																
<i>Eusarsiella (LPIL)</i>																																
<i>Eusarsiella radicosta</i>																																
<i>Eusarsiella texana</i>	1	.25																														

**Table D-2. (continued)**

Species List	D1		D2		D3		D4		Mean D		E1		E2		E3		E4		Mean E		F1		F2		F3		F4		Mean F	
	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density		
Order PODOCOPIDA <i>Podocopida (LPIL)</i>																														
<b>BRYOZOA</b> <i>Bryozoa (LPIL)</i>																														
<b>CHORDATA</b>																														
CLASS LEPTOCARDIA																														
Order AMPHIOXI																														
FAMILY BRANCHIOSTOMIDAE																														
<i>Branchiostoma (LPIL)</i>																												.75 18.75		
<b>CNIDARIA</b>																														
CLASS HYDROZOA																														
<i>Hydrozoa (LPIL)</i>																														
<b>ECHINODERMATA</b>																														
<i>Echinodermata (LPIL)</i>																														
CLASS ECHINOIDEA																														
<i>Echinoidea (LPIL)</i>																														
Order CLYPEASTEROIDA																														
FAMILY MELLITIDAE																														
<i>Mellitidae (LPIL)</i>																												.50 .50 12.50		
<i>Mellita isometra</i>																														
CLASS OPHIUROIDEA																														
<i>Ophiuroidea (LPIL)</i>																												.50 12.50		
Order OPHIURIDA																														
FAMILY AMPHIURIDAE																														
<i>Amphiuridae (LPIL)</i>																														
<b>MOLLUSCA</b>																														
CLASS BIVALVIA																														
<i>Bivalvia (LPIL)</i>																														
Order OSTREOIDA																														
FAMILY ANOMIIDAE																														
<i>Anomia simplex</i>																												.25 6.25		
Order VENEROIDA																														
FAMILY CRASSATELLIDAE																														
<i>Crassinella dupliniana</i>																												1.00 25.00		
<i>Crassinella lunulata</i>																												.75 18.75		
FAMILY LUCINIDAE																														
<i>Lucinidae (LPIL)</i>	1	25																										.75 18.75		
<i>Lucina (LPIL)</i>																														
<i>Lucina multilineata</i>																														
FAMILY MESODESMATIDAE																														
<i>Ervilia concentrica</i>																														

**Table D-2. (concluded)**

Species List	D1				D2				D3				D4				Mean D		E1				E2				E3				E4				Mean E		F1				F2				F3				F4				Mean F	
	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density	Count	Density																										
FAMILY MONTACUTIDAE																																																						
<i>Montacutidae (LPIL)</i>																																																						
FAMILY SEMELIDAE																																																						
<i>Semelidae (LPIL)</i>																																																						
FAMILY TELLINIDAE																																																						
<i>Strigilla mirabilis</i>																																																						
<i>Tellina (LPIL)</i>																																																						
<i>Tellina iris</i>																																																						
CLASS GASTROPODA																																																						
Order CEPHALASPIDEA																																																						
FAMILY SCAPHANDRIDAE																																																						
<i>Acteocina canaliculata</i>																																																						
<i>Cylichna alba</i>	1	25																																																				
Order MESOGASTROPODA																																																						
FAMILY CAECIDAE																																																						
<i>Caecum pulchellum</i>			1	25	3	75	3	75	1.75	43.75	3	75	7	175	12	300	2	50	6.00	150.00																																		
FAMILY EULIMIDAE																																																						
<i>Strombiformis bilineatus</i>																																																						
FAMILY NATICIDAE																																																						
<i>Tectonatica pusilla</i>																																																						
Order NEOGASTROPODA																																																						
FAMILY OLIVIDAE																																																						
<i>Oliva sayana</i>																																																						
<i>Olivella dealbata</i>																																																						
FAMILY TURRIDAE																																																						
<i>Kurtziella astrostyla</i>																																																						
Order PYRAMIDELLOIDA																																																						
FAMILY PYRAMIDELLIDAE																																																						
<i>Odostomia weberi</i>																																																						
<i>Turbonilla (LPIL)</i>																																																						
PHORONIDA																																																						
FAMILY PHORONIDAE																																																						
<i>Phoronis (LPIL)</i>																																																						
PLATYHELMINTHES																																																						
CLASS TURBELLARIA																																																						
<i>Turbellaria (LPIL)</i>																																																						
RHYNCHOCOELA																																																						
<i>Rhynchocoela (LPIL)</i>	1	25																																																				
SIPUNCULA																																																						
FAMILY ASPIDOSIPHONIDAE																																																						
<i>Aspidosiphon albus</i>																																																						

**Table D-3. Biomass for major taxa groups from Topsail Beach, NC, benthic sampling stations.**

<b>Station: A1</b>	<b>gm/04 m<sup>2</sup></b>	<b>gm/m<sup>2</sup></b>
Annelida	0.5556	13.89
Mollusca	0.5544	13.86
Arthropoda	0.0000	0
Echinodermata	0.0000	0
Other Taxa	0.0000	0
<b>Total</b>	<b>1.1100</b>	<b>27.75</b>

<b>Station: A2</b>	<b>gm/04 m<sup>2</sup></b>	<b>gm/m<sup>2</sup></b>
Annelida	0.6517	16.2925
Mollusca	0.5792	14.48
Arthropoda	0.0000	0
Echinodermata	0.0000	0
Other Taxa	0.0000	0
<b>Total</b>	<b>1.2309</b>	<b>30.7725</b>

<b>Station: A3</b>	<b>gm/04 m<sup>2</sup></b>	<b>gm/m<sup>2</sup></b>
Annelida	0.5557	13.8925
Mollusca	0.5512	13.78
Arthropoda	0.5484	13.71
Echinodermata	0.0000	0
Other Taxa	0.5490	13.725
<b>Total</b>	<b>2.2043</b>	<b>55.1075</b>

<b>Station: A4</b>	<b>gm/04 m<sup>2</sup></b>	<b>gm/m<sup>2</sup></b>
Annelida	0.5620	14.05
Mollusca	0.5712	14.28
Arthropoda	0.5563	13.9075
Echinodermata	0.0000	0
Other Taxa	0.0000	0
<b>Total</b>	<b>1.6895</b>	<b>42.2375</b>

<b>Station: A5</b>	<b>gm/04 m<sup>2</sup></b>	<b>gm/m<sup>2</sup></b>
Annelida	0.5517	13.7925
Mollusca	0.5522	13.805
Arthropoda	0.0000	0
Echinodermata	0.0000	0
Other Taxa	0.0000	0
<b>Total</b>	<b>1.1039</b>	<b>27.5975</b>

<b>Station: B1</b>	<b>gm/04 m<sup>2</sup></b>	<b>gm/m<sup>2</sup></b>
Annelida	0.5519	13.7975
Mollusca	0.5513	13.7825
Arthropoda	0.5711	14.2775
Echinodermata	0.5760	14.4
Other Taxa	0.5654	14.135
<b>Total</b>	<b>2.8157</b>	<b>70.3925</b>

<b>Station: B2</b>	<b>gm/04 m<sup>2</sup></b>	<b>gm/m<sup>2</sup></b>
Annelida	0.5520	13.8
Mollusca	0.5520	13.8
Arthropoda	0.5481	13.7025
Echinodermata	0.5492	13.73
Other Taxa	0.5491	13.7275
<b>Total</b>	<b>2.7504</b>	<b>68.76</b>

<b>Station: B3</b>	<b>gm/04 m<sup>2</sup></b>	<b>gm/m<sup>2</sup></b>
Annelida	0.5665	14.1625
Mollusca	0.5511	13.7775
Arthropoda	0.5513	13.7825
Echinodermata	0.5490	13.725
Other Taxa	0.5595	13.9875
<b>Total</b>	<b>2.7774</b>	<b>69.435</b>

<b>Station: C1</b>	<b>gm/04 m<sup>2</sup></b>	<b>gm/m<sup>2</sup></b>
Annelida	0.5526	13.815
Mollusca	0.5490	13.725
Arthropoda	0.5520	13.8
Echinodermata	0.0000	0
Other Taxa	0.0000	0
<b>Total</b>	<b>1.6536</b>	<b>41.34</b>

<b>Station: C2</b>	<b>gm/04 m<sup>2</sup></b>	<b>gm/m<sup>2</sup></b>
Annelida	0.5526	13.815
Mollusca	0.5490	13.725
Arthropoda	0.5520	13.8
Echinodermata	0.0000	0
Other Taxa	0.0000	0
<b>Total</b>	<b>1.6536</b>	<b>41.34</b>

<b>Station: C3</b>	<b>gm/04 m<sup>2</sup></b>	<b>gm/m<sup>2</sup></b>
Annelida	0.5556	13.89
Mollusca	0.5555	13.8875
Arthropoda	0.5564	13.91
Echinodermata	0.5491	13.7275
Other Taxa	0.5496	13.74
<b>Total</b>	<b>2.7662</b>	<b>69.155</b>

<b>Station: C4</b>	<b>gm/04 m<sup>2</sup></b>	<b>gm/m<sup>2</sup></b>
Annelida	0.5539	13.8475
Mollusca	0.5496	13.74
Arthropoda	0.5533	13.8325
Echinodermata	21.4873	537.1825
Other Taxa	0.0000	0
<b>Total</b>	<b>23.1441</b>	<b>578.6025</b>

**Table D-3. (concluded)**

Station: D1	gm/04 m <sup>2</sup>	gm/m <sup>2</sup>	Station: E2	gm/04 m <sup>2</sup>	gm/m <sup>2</sup>
Annelida	0.5556	13.89	Annelida	0.5546	13.865
Mollusca	0.5494	13.735	Mollusca	0.5545	13.8625
Arthropoda	0.5483	13.7075	Arthropoda	0.5563	13.9075
Echinodermata	0.0000	0	Echinodermata	0.5492	13.73
Other Taxa	0.0000	0	Other Taxa	0.5490	13.725
Total	1.6533	41.3325	Total	2.7636	69.09

Station: D2	gm/04 m <sup>2</sup>	gm/m <sup>2</sup>	Station: E3	gm/04 m <sup>2</sup>	gm/m <sup>2</sup>
Annelida	0.5659	14.1475	Annelida	0.5622	14.055
Mollusca	0.5491	13.7275	Mollusca	0.5522	13.805
Arthropoda	0.0000	0	Arthropoda	0.5563	13.9075
Echinodermata	0.0000	0	Echinodermata	0.5491	13.7275
Other Taxa	0.5492	13.73	Other Taxa	0.5490	13.725
Total	1.6642	41.605	Total	2.7688	69.22

Station: D3	gm/04 m <sup>2</sup>	gm/m <sup>2</sup>	Station: E4	gm/04 m <sup>2</sup>	gm/m <sup>2</sup>
Annelida	0.5658	14.145	Annelida	0.5623	14.0575
Mollusca	0.5493	13.7325	Mollusca	0.5543	13.8575
Arthropoda	0.0000	0	Arthropoda	0.5524	13.81
Echinodermata	0.0000	0	Echinodermata	0.0000	0
Other Taxa	0.0000	0	Other Taxa	0.5492	13.73
Total	1.1151	27.8775	Total	2.2182	55.455

Station: D4	gm/04 m <sup>2</sup>	gm/m <sup>2</sup>	Station: F2	gm/04 m <sup>2</sup>	gm/m <sup>2</sup>
Annelida	0.5743	14.3575	Annelida	0.6508	16.27
Mollusca	0.5495	13.7375	Mollusca	0.5512	13.78
Arthropoda	0.5485	13.7125	Arthropoda	0.5561	13.9025
Echinodermata	0.5492	13.73	Echinodermata	0.0000	0
Other Taxa	0.0000	0	Other Taxa	0.6559	16.3975
Total	2.2215	55.5375	Total	2.4140	60.35

Station: E1	gm/04 m <sup>2</sup>	gm/m <sup>2</sup>	Station: F3	gm/04 m <sup>2</sup>	gm/m <sup>2</sup>
Annelida	0.5722	14.305	Annelida	0.6508	16.27
Mollusca	0.5526	13.815	Mollusca	0.5512	13.78
Arthropoda	0.5766	14.415	Arthropoda	0.5561	13.9025
Echinodermata	0.0000	0	Echinodermata	0.0000	0
Other Taxa	0.0000	0	Other Taxa	0.6559	16.3975
Total	1.7014	42.535	Total	2.4140	60.35

Station: F1	gm/04 m <sup>2</sup>	gm/m <sup>2</sup>	Station: F4	gm/04 m <sup>2</sup>	gm/m <sup>2</sup>
Annelida	0.5520	13.8	Annelida	0.5513	13.7825
Mollusca	0.5513	13.7825	Mollusca	0.5740	14.35
Arthropoda	0.5520	13.8	Arthropoda	0.5490	13.725
Echinodermata	0.0000	0	Echinodermata	0.5496	13.74
Other Taxa	0.5491	13.7275	Other Taxa	0.0000	0
Total	2.2044	55.11	Total	2.2239	55.5975